## STAC Workshop Climate Change Modeling 2.0

Gary Shenk CBPO

MWG

October 10, 2021

#### STAC Climate Change Modeling 2.0

- Workshop held in 2018 in response to PSC decision in 2017 to take a longer look at climate change as part of the Midpoint Assessment
- Purpose: Guidance on the models and assessment framework used to assess the effect of climate change on the TMDL

#### **Workshop Steering Committee:**

Co-Chair: Mark Bennett (USGS) Co-Chair: Lewis Linker (EPA CBPO)

Don Boesch (UMCES)

Lee Currey (MD Dept. of Environment)

Marjorie Friedrichs (VIMS)

Maria Herrmann (Penn State) \*

Raleigh Hood (UMCES)

Tom Johnson (USEPA)\*

Andy Miller (UMBC)\*

Dave Montali (Tetra Tech)

**STAC Staff:** 

Rachel Dixon, Chesapeake Research Consortium Annabelle Harvey, Chesapeake Research Consortium


Chesapeake Bay Program **Climate Change Modeling 2.0** 



STAC Workshop Report September 24-25, 2018 Annapolis, MD

STAC Publication 21-003

#### Why is it coming out now?




Too busy implementing recommendations!

#### Near Term Recommendations (by 2021)

- Introduce the estimation of uncertainty into the decision process.
- Closely **examine** all climate-related sensitivities in the watershed and estuarine models and characterize the relative importance of these effects on dissolved oxygen.
- With the knowledge gained from sensitivity analyses of recommendation #2, prioritize **improvements** and enhanced quantification of uncertainties in model inputs.
- Compare **multiple** existing estuarine **model** simulations, including the CBP's Water Quality and Sediment Transport Model, from 1985 through recent wet years.

# Near term: Closely examine sensitivities and make improvements



## Near term: Closely examine sensitivities and make improvements- watershed

- Spatially vary the relationship between nitrogen and flow
- Developed area sensitivity to Phosphorus
- Investigate potential changes to wastewater overflows
- Change the nitrogen and phosphorus speciation methods
- Incorporate the effects of climate change into the BMP effectiveness estimates

Climate change documentation: https://cast.chesapeakebay.net/FileBrowser/GetFile?fileName= P6ModelDocumentation%2FClimateChangeDocumentation.pdf


- Incorporate the effects of climate change into the BMP effectiveness estimates
  - Virginia Tech scientists are producing a comprehensive literature survey on the effects of climate change on nutrient cycling and BMP effectiveness.
  - WQGIT-funded project produced <u>updated intensity-duration-frequency curves</u> for all counties in the watershed will supply necessary estimates of climate effects.

- Resolve the effect of sea-level rise
- Increase phytoplankton growth at high temperatures.
- Include a temperature dependent mortality or grazing rate in the model.
  - Re-examine the outer boundary conditions
  - Investigate downscaled future wind products.

Near Term: Compare Multiple Water Quality

Models

- VIMS-led CHAMP project
  - Multiple climate models
  - Multiple watershed models
  - Multiple estuarine models



direct links between various components of the ecosystembased scenario forecast modeling system.

#### Near Term Recommendations (by 2021)



• Introduce the estimation of **uncertainty** into the decision process.



 Closely examine all climate-related sensitivities in the watershed and estuarine models and characterize the relative importance of these effects on dissolved oxygen.



• With the knowledge gained from sensitivity analyses of recommendation #2, prioritize **improvements** and enhanced quantification of uncertainties in model inputs.



 Compare multiple existing estuarine model simulations, including the CBP's Water Quality and Sediment Transport Model, from 1985 through recent wet years.

#### In the longer term (by 2025)


- Develop a new estuarine model
- Continue development of climate-related watershed model capabilities with particular attention to BMP effectiveness.
- Create a more **sophisticated evaluation framework** that incorporates various sources of uncertainty:

#### Long Term: Develop a new estuarine model

- Use an unstructured grid
- Simulate wetting and drying
- Develop a spectral wave model
- Extend the boundary onto the coastal shelf
- Improve biogeochemical parameterizations
- Improve model forcing

### Long Term: Estuarine model RFA

- Unstructured Grid
- May be part of new model
  - Simulate wetting and drying
  - Develop a spectral wave model
  - Extend the boundary onto the coastal shelf
  - Improve biogeochemical parameterizations
  - Improve model forcing



#### Long Term: Watershed model

 Continue watershed model refinement with particular attention to BMP effectiveness


- Modeling team working on scale issues
- Waiting on prioritization from WQGIT and other partners

- Climate inputs
  - Use models rather than trend
  - Evaluate multiple RCPs (now SSPs)
  - Use climate models at a finer spatial scale
- Develop improved suite of mechanistic models
- Interact with similar efforts world-wide
- Create a climate research strategy
- Re-evaluate goals relative to climate change
- Use uncertainty in the decision process


- Climate inputs
  - Use climate models rather than observed trend
  - Evaluate multiple RCPs (now SSPs)
  - Use climate models at a finer spatial scale
- CBP will consider these in 2024-2025
- Develop improved suite of mechanistic models
- Yes for estuarine model: No for watershed model
- Interact with similar efforts world-wide
- Yes for scientists; ??? for managers

- Create a climate research strategy
  - the CBP should consider developing, supporting, and periodically updating a Strategic Research Framework that identifies ongoing activities and priority knowledge gaps.


- Create a climate research strategy
  - the CBP should consider developing, supporting, and periodically updating a Strategic Research Framework that identifies ongoing activities and priority knowledge gaps.



- Re-evaluate goals relative to climate change
  - CBP noted issues with the Open Water designated uses under climate change and declined to base reductions on Open Water
  - Criteria Assessment Protocols Workgroup looking at changing Open Water boundaries.
  - Potential issues will arise with assessment shortduration Water Quality Standards



#### Use of uncertainty in Decision Process




- Workshop Suggestion: Use of multiple scenarios to define a probability of attainment for a particular load reduction goal.
- CBP could consider in 2024-2025


#### Summary


- Watershed modeling
  - Significant advances incorporated into 2020 climate decision
  - Little additional work specified
- Estuarine modeling
  - Significant advances incorporated into 2020 climate decision
  - Progress being made toward wholesale changes recommended for the future
- Overall framework
  - Challenge to incorporate uncertainty in decision-making
  - Re-examine climate assumptions
  - Re-examine environmental endpoints

### Extra Slides

Spatially vary the relationship between nitrogen and flow









No consistent relationship to land use found


- Spatially vary the relationship between nitrogen and flow
- Using monitoring data, a slightly higher than a 1:1 relationship between flow change and TN change found in lower loading areas, but not a strong enough relationship to include.



Developed area sensitivity to Phosphorus

#### Literature review shows 1:1 relationship to flow







# CSO volume


## Near term: Closely examine sensitivities - watershed

Investigate potential changes to wastewater overflows






Change the nitrogen and phosphorus speciation methods




- Nitrate fraction increases with more load across space
- Nitrate fraction decreases with more load at a single station.


Resolve the effect of sea-level rise



Increase phytoplankton growth at high temperatures.




 Include a temperature dependent mortality or grazing rate in the model.




 More recent and local literature used to modify respiration curve Near term: Closely examine sensitivities -

estuarine

- Re-examine the outer boundary conditions
- Change in surface temperature is 90% of the air temperature change
- Change in temperature at depth is the surface change times the ratio of initial temperatures – colder, deeper water changes less



Investigate downscaled future wind products.




 Need around 4 meters per second to make a significant difference

 Change in wind speed on the order of 0.1 meters per second

#### Near term: Uncertainty


 Evaluated uncertainty of rainfall and temperature extrapolations



#### Near term: Uncertainty

Evaluated uncertainty of individual components


 But did not put it all together to estimate overall uncertainty



Near Term: Compare Multiple Water Quality

Models

- VIMS-led CHAMP project
  - Multiple climate models
  - Multiple watershed models
  - Multiple estuarine models



direct links between various components of the ecosystembased scenario forecast modeling system.