


# Generalized Additive Model Development for Patuxent Horizontal Chlorophyll.

Elgin Perry  
TMAW  
Dec. 4, 2013

# GAM project overview

- Developed gams for Patuxent Horizontal Chla
- Model development examined season, flow, light and nonlinear trend.
- New tool for season using cyclical day of year.
- Thin plate splines for modeling changes in the flow to response relationship over time.


# Trend(season,flow)




# Steps

1. Look at response data
2. Look at data relative to forcing functions
3. Develop quantitative measures of forcing functions
4. GAM development.


# Response data (388 dates)




# Late Summer



# Late Summer




# Late Summer







# Spring Bloom




# Spring Bloom




# Spring Bloom




# Spring Bloom




# Early Winter




# Early Winter



# Early Winter




# Early Winter







# Mid-late Winter




# Mid-late Winter




# Mid-late Winter




# Mid-late Winter






# Mid-late Winter




# Spring Bloom














# Patterns

High flow reduces Chl in TF.

High Chl in TF sets up during low flow periods

Flow appears to have little effect downriver

Flow and light are connected


Maybe cloudy days lead to lower Chl.


# Quantitative Measures


- Dependent – Integral of chl by salinity zone
- Season
  - Monthly means
  - Cyclical day of year (doy)
- Flow
  - Moving average of flow with lags (up to 7 weeks)
  - Days since flow event
  - Size of flow event
- Light
  - Series of averages ( 1 day, 2 day, . . . 7 day)

# Quantitative Measures


- Light
  - Series of averages ( 1 day, 2 day, . . . 7 day)
  - Sunny days ( $\text{par} > 0.75$  quantile)
  - Cloudy days ( $\text{par} < 0.25$  quantile)
  - Sunny days-cloudy days




















































# Variable Added Plots












# Finally a gam trend line



# Seasonally adjusted trend



# Trend(season,flow)



# Conclusions

TF chlorophyll is going down

Effective management actions were in 1980's and post 2000.

# Analysis Conclusions

- Cyclical DOY is good seasonal method
- No season x time interaction
- No flow x time interaction
- Event.days, Event.size = deadend
- Light = deadend

