


A faint, light blue map of the Patuxent River estuary is visible in the background. The map shows the river's complex, branching structure as it flows into the Chesapeake Bay. The land areas are depicted in a light yellowish-green color.

Patuxent River Estuary Model

Progress in SCHISM–ICM Development and Validation



Amir Reza Azarnivand and Jeremy M. Testa

Chesapeake Biological Lab, University of Maryland Center for Environmental Science

10/28/2026

From Hydrodynamics to Water Quality

- 3-year sensitivity runs (1991-1993) to make decisions about the best I.C. for the longer run.
- Only ICM (NO external sediment model)
- 3 different sets of data in this period: CBP fixed stations, Solomons Harbor stations, Boynton et al. flux data
- Jeremy Testa's stand-alone SFM work on Patuxent stations used for I.C. (Sediment POC, PON, POP, NH4 and PO4)



Three Challenges in Model Development

1) P sorption: How sensitive are phosphorus fluxes to initial sediment conditions?

Tests:

v1: bNH40 = 4.0, bNO30 = 1.0, bPO40 = 5.0, bpiePO4 = 180.0, bKOPO4f = 3000.0, bKOPO4s = 300.0

v5: bNH40 = -999, bNO30 = -999, bPO40 = -999, bpiePO4 = 10.0, bKOPO4f = 700.0, bKOPO4s = 150.0

POC(x9), PON(x3), POP(x3 to G1 and G3, a lot more to G2)

v6: bNH40 = -999, bNO30 = -999, bPO40 = -999, bpiePO4 = 100.0, bKOPO4f = 700.0, bKOPO4s = 150.0

POC(x9), PON(x3), POP(x3 to G1 and G3, a lot more to G2)

2) Sediment oxygen demand and bottom DO: How do initial sediment conditions affect sediment oxygen demand and bottom-water oxygen?

Tests:

v1: bNH40 = 4.0, bNO30 = 1.0, bPO40 = 5.0, bpiePO4 = 180.0, bKOPO4f = 3000.0, bKOPO4s = 300.0

v5: bNH40 = -999, bNO30 = -999, bPO40 = -999, bpiePO4 = 10.0, bKOPO4f = 700.0, bKOPO4s = 150.0

POC(x9), PON(x3), POP(x3 to G1 and G3, a lot more to G2)

v6: bNH40 = -999, bNO30 = -999, bPO40 = -999, bpiePO4 = 100.0, bKOPO4f = 700.0, bKOPO4s = 150.0

POC(x9), PON(x3), POP(x3 to G1 and G3, a lot more to G2)



3) Solomons Harbor water-level issue

Tests:



Distributed freshwater sources, reduced dt, increased minimum water depth for wetting/drying

1) P sorption – Sediment PP

Sediment particulate organic P: model vs observations (1991–1993)





1) P sorption – Sediment PP





1) P sorption – Sediment PO_4 Flux

Sediment PO_4 Flux: Model vs. Observations (1991–1993)





1) P sorption – Sediment PO_4 Flux





2) Sediment oxygen demand

Sediment oxygen demand — Benthic stations (1991–1993)





2) Sediment oxygen demand





2) Sediment oxygen demand

Sediment oxygen demand — Solomons Harbor (1991–1993)





2) Bottom DO





2) Bottom DO

Bottom DO | Main channel | 1991-1993




2) Bottom DO




3) Solomons Harbor Water Level – in progress

Before





$dt = 150$
 $h0 = 5 \text{ cm}$

After





3) Solomons Harbor Water Level – in progress

Water level in Solomons Harbor (1991–1993)






Solomons Harbor

Surface Temperature



Bottom Salinity

Surface Chl-a





Thank you