

**Recommended relative-to-rye phosphorus reduction efficiencies (RPRE) and relative-to-rye sediment reduction efficiencies (RSRE) based on best professional judgment of panel members.**

Panelist	Annual Ryegrass		Annual Legume		Legume + Grass Mix.		Brassica (winter hardy)		Forage Radish		Forage Radish + Grass Mix.		Triticale		Oats (winter hardy)		Oats (winter killed)	
	RPRE	RSRE	RPRE	RSRE	RPRE	RSRE	RPRE	RSRE	RPRE	RSRE	RPRE	RSRE	RPRE	RSRE	RPRE	RSRE	RPRE	RSRE
Anonymous A	0.6	0.7	0.2	0.5	0.4	0.4	0.5	0.6	0.6	0.6	0.6	0.6	0.5	0.9	0.4	0.4	0.3	0.3
Anonymous B	1.0	0.9	1.0	0.7	1.1	1.0	1.2	0.9	0.6	0.7	0.9	1.0	1.0	0.9	0.9	0.9	0.8	0.8
Anonymous C	0.7	0.7	0.2	0.3	0.6	0.6	0.6	0.6	0.4	0.5	0.5	0.5	0.9	0.9	0.3	0.4	0.3	0.4
Anonymous D	1.0	1.0	0.5	0.5	1.0	1.0	0.7	0.7	0.6	0.6	0.6	0.6	0.9	1.0	0.7	0.9	0.4	0.4
Anonymous E	0.7	0.8	0.5	0.8	0.6	0.8	0.7	0.8	0.6	0.6	0.6	0.8	0.9	0.8	0.6	0.7	0.4	0.4
Anonymous F	0.7	0.8	0.5	0.4	0.6	0.7	0.7	0.4	0.6	0.3	0.6	0.5	0.9	0.9	0.8	0.9	0.4	0.5
Anonymous G	0.8	0.8	0.2	0.2	0.5	1.0	0.5	0.7	0.0	0.4	0.2	0.7	1.0	1.0	1.0	1.0	0.2	0.2
Anonymous H	0.7	0.7	0.2	0.2	0.4	0.4	0.7	0.7	0.3	0.3	0.5	0.5	0.9	0.9	0.6	0.6	0.3	0.3
Anonymous I	0.3	0.3	0.1	0.1			0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3		
Anonymous J	0.7	0.8	0.4	0.6	0.7	0.8	0.5	0.7	0.0	0.1	0.4	0.5	1.0	1.0	0.7	0.8	0.2	0.4
<b>Average</b>	<b>0.70</b>	<b>0.73</b>	<b>0.38</b>	<b>0.42</b>	<b>0.66</b>	<b>0.75</b>	<b>0.65</b>	<b>0.65</b>	<b>0.40</b>	<b>0.44</b>	<b>0.54</b>	<b>0.62</b>	<b>0.81</b>	<b>0.85</b>	<b>0.63</b>	<b>0.68</b>	<b>0.37</b>	<b>0.41</b>
<i>Standard Error</i>	<i>0.06</i>	<i>0.06</i>	<i>0.08</i>	<i>0.07</i>	<i>0.08</i>	<i>0.08</i>	<i>0.07</i>	<i>0.05</i>	<i>0.08</i>	<i>0.06</i>	<i>0.06</i>	<i>0.06</i>	<i>0.07</i>	<i>0.07</i>	<i>0.07</i>	<i>0.08</i>	<i>0.06</i>	<i>0.06</i>
Average with high and low	0.68	0.76	0.33	0.41	0.66	0.71	0.61	0.69	0.35	0.45	0.54	0.60	0.81	0.86	0.66	0.69	0.33	0.39
Median	0.68	0.80	0.30	0.45	0.60	0.80	0.65	0.70	0.48	0.43	0.55	0.57	0.86	0.88	0.63	0.75	0.33	0.40