# The Water Quality Standards Indicator and its Friends

Peter Tango USGS@CBPO
Chesapeake Bay Monitoring Coordinator
STAR Meeting
June 26, 2025

### Reminder – The WQSAM Outcome (2014)

Continually improve the capacity to monitor and assess the effects
of management actions being undertaken to implement the Bay
TMDL and improve water quality. Use the monitoring results to report
annually to the public on progress made in attaining established Bay
water-quality standards and trends in reducing nutrients and
sediment in the watershed.

### Reminder – The WQSAM Outcome (2014)

• Continually improve the capacity to monitor and assess the effects of management actions being undertaken to implement the Bay TMDL and improve water quality. Use the monitoring results to report annually to the public on progress made in attaining established Bay water-quality standards and trends in reducing nutrients and sediment in the watershed.




Exciting?



Compelling?


Happy marriage in creating the indicator Modeling scenario
 "stoplight plots" of the
 TMDL accounting for
 meeting water quality
 standards





Conditional Probability assessment for gap filling monitoring results:

- USEPA 2004
- USEPA 2017



# Peer reviewed foundations behind the WQSAM indicator



#### Science of The Total Environment


Volumes 637-638, 1 October 2018, Pages 1617-1625




Short Communication

Chesapeake Bay's water quality condition has been recovering: Insights from a multimetric indicator assessment of thirty years of tidal monitoring data

Qian Zhang <sup>a</sup>  $\stackrel{\triangle}{\sim}$   $\stackrel{\boxtimes}{\bowtie}$ , Rebecca R. Murphy <sup>a</sup>, Richard Tian <sup>a</sup>, Melinda K. Forsyth <sup>b</sup>, Emily M. Trentacoste <sup>c</sup>, Jennifer Keisman <sup>d</sup>, Peter J. Tango <sup>e</sup>





# Mythbuster Metrics: Beyond the WQSAM Indicator...

# Dissolved oxygen (mg L<sup>-1</sup>) s02 (Poor) 02 to 1 1 to 2 2 to 3 3 to 5 > 5 (Good) Richmond Richmond Norfolk





Figure 4: Occurrences of hypoxia and anoxia are higher in the summer months than the winter months because oxygen-consuming processes increase with temperature. Data from 1985–2005.

# Beyond WQ Stds attainment – diverse, important habitat metrics

- Dead zone length
- Dead zone area
- Dead zone volume
- Start date, end date
- Speed of hypoxia formation
- Duration
- Intensity
- Attainment deficit, attainment buffer

Who cares about length of hypoxia in the bay?



Our scientists do @...

#### **Ecological Forecasting and** the Science of Hypoxia in **Chesapeake Bay**

CLARK, WILLIAM C. DENNISON, E. CAROLINE DONOVAN, ALEXANDER W. FISHER,

...because it is one the foundations to how we forecast summer hypoxia for Chesapeake Bay

### University of Michigan Chesapeake Bay hypoxia forecasting model

#### Driver:

Jan-May average Susquehanna TN load



#### Calibration target:

Mean July hypoxic volume (HV) ([DO] < 2 mg/L)




#### Model output:

Average subpycnocline [DO] as a function of distance from TN source



**Hypoxic length** = sum of all segments with [DO] < 2 mg/L



Hypoxic length → hypoxic volume through empirical V-L relationship

# Length of bay with hypoxia also tells a story of recovery, aligns with our science understanding of bay behavior through time



Contents lists available at ScienceDirect

#### Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv



Nitrogen reductions have decreased hypoxia in the Chesapeake Bay: Evidence from empirical and numerical modeling




Luke T. Frankel <sup>a,\*</sup>, Marjorie A.M. Friedrichs <sup>a</sup>, Pierre St-Laurent <sup>a</sup>, Aaron J. Bever <sup>b</sup>, Romuald N. Lipcius <sup>a</sup>, Gopal Bhatt <sup>c,d</sup>, Gary W. Shenk <sup>c,e</sup>

- <sup>a</sup> Virginia Institute of Marine Science, William & Mary, 1370 Greate Road, Gloucester Point, VA, USA
- b Anchor QEA LLC, 1201 3rd Avenue, Suite 2600, Seattle, WA, USA
- <sup>c</sup> Chesapeake Bay Program Office, 1750 Forest Drive, Suite 130, Annapolis, MD, USA
- d Department of Civil & Environmental Engineering, The Pennsylvania State University, 212 Sackett Building, University Park, PA, USA
- <sup>e</sup> U.S. Geological Survey, Virginia and West Virginia Water Science Center, 1730 East Parham Road, Richmond, VA, USA

Frankel et al. 2022

#### GRAPHICAL ABSTRACT



Myth: The Summer Dead Zone is June-September

The dead zone can start in March and end in November






Figure 4: Occurrences of hypoxia and anoxia are higher in the summer months than the winter months because oxygen-consuming processes increase with temperature. Data from 1985–2005.

**UMCES-Ecocheck 2007** 

\* Tracking starting dates and ending dates are important phenological metrics for informing and explaining system changes

## Seasonal timing (phenology) affects biology







# Myth: July is the worst month for the Dead Zone

# Myth: July is the worst month




Figure 4: Occurrences of hypoxia and anoxia are higher in the summer months than the winter months because oxygen-consuming processes increase with temperature. Data from 1985–2005.

UMCES-Ecocheck 2007

There are variations on the expression of **peak timing.**Some such metrics have shown shifts in hypoxia timing

### Long-Term Trends in Chesapeake Bay Seasonal Hypoxia, Stratification, and Nutrient Loading

Rebecca R. Murphy, W. Michael Kemp and William P. Ball

**Estuaries and Coasts** 

Vol. 34, No. 6 (NOVEMBER 2011), pp. 1293-1309 (17 pages)


Published By: Springer Nature

# Myth: Fish die in the Dead Zone


Tidal Bay Monitoring. Degraded deep water habitats, however,

fish kills and algal blooms are reported annually throughout

shallow waters of the estuary.



Management and monitoring target mismatch?











Tidal Bay Monitoring. Degraded deep water habitats, however, fish kills and algal blooms are reported annually throughout


shallow waters of the estuary. Achieving Water Quality Goals in the Chesapeake Bay: A Comprehensive Evaluation of System Response AS EXAMINATION OF ROW THE CHESAPEANX BAY ECOSYSTEM HAS RESPONDED.





...fix this?








# September 26, 2005 Across the water...MD DNR investigates a local fish kill













#### September 28, 2005 Estimated 30,000-50,000 fish dead; 15 species affected







Fish Kill in the Corsica



## Building Environmental Intelligence 4. Fish necropsy work for toxin-specific effects





# Myth: There is one Dead Zone in the bay

### Patuxent River has one

Estuaries Vol. 26, No. 2A, p. 280-297 April 2003

# The Pattern and Influence of Low Dissolved Oxygen in the Patuxent River, a Seasonally Hypoxic Estuary

DENISE L. BREITBURG<sup>1,\*</sup>, AARON ADAMACK<sup>2</sup>, KENNETH A. ROSE<sup>2</sup>, SARAH E. KOLESAR<sup>1</sup>, MARY BETH DECKER<sup>1,3,†</sup>, JENNIFER E. PURCELL<sup>3,‡</sup>, JULIE E. KEISTER<sup>4</sup>, and JAMES H. COWAN, JR.<sup>2</sup>


<sup>&</sup>lt;sup>1</sup> The Academy of Natural Sciences, Estuarine Research Center, 10545 Mackall Road, St. Leonard, Maryland 20678


<sup>&</sup>lt;sup>2</sup> Coastal Fisheries Institute and Department of Oceanography and Coastal Sciences, Wetland Resources Building, Louisiana State University, Baton Rouge, Louisiana 70803

<sup>&</sup>lt;sup>3</sup> Horn Point Laboratory, University of Maryland Center for Environmental Science, Cambridge, Maryland 2113-0775

<sup>&</sup>lt;sup>4</sup> Hatfield Marine Science Center, Oregon State University, 2030 South Marine Science Drive, Newport, Oregon 97365

### Severn River dead zone





Article

PDF Available

Resolving spatiotemporal characteristics of the seasonal hypoxia cycle in shallow estuarine environments of the Severn River and South River, MD, Chesapeake Bay, USA

September 2016 · Heliyon 2(9):e00157

DOI:10.1016/j.heliyon.2016.e00157

License · CC BY 4.0

Authors:



Andrew Muller
United States Naval Academy





Diana L Muller



Arianna Muller

Geographic distribution of hypoxia could be improved in our graphics and communications with such indicators: touch the local scale





> Environ Sci Technol. 2025 Jan 14;59(1):553-564. doi: 10.1021/acs.est.4c07368. Epub 2024 Dec 26.

Geography, Trajectories, and Controls of Coastal Water Quality: More Rapid Improvement in the Shallow Zone of the Chesapeake Bay


Qian Zhang <sup>1</sup>, Rebecca R Murphy <sup>1</sup>, Richard Tian <sup>1</sup>, Peter J Tango <sup>2</sup>

Affiliations + expand

PMID: 39723897 PMCID: PMC11740901 (available on 2025-12-26) DOI: 10.1021/acs.est.4c07368

Myth: Water Quality
Standards is the best
way to track change in
the bay

# Hypoxic Volume Days is a more management responsive metric with our data set than WQS attainment, July maximum hypoxia




The pace of recovery over 30 years was measurable in hypoxia duration;

The pace was about 0.8 days hypoxia improvement per year.

# We also have detailed evaluations of bay conditions from Attainment deficits and Attainment buffers from our WQSAM analysis

### **ORIGINAL RESEARCH article** This article is part of the Research Topic Front. Mar. Sci., 20 November 2018 Research and Management of Sec. Marine Ecosystem Ecology **Eutrophication in Coastal Ecosystems** Volume 5 - 2018 | https://doi.org/10.3389/fmars.2018.00422 View all 23 articles > Chesapeake Bay Dissolved Oxygen Criterion Attainment Deficit: Three Decades of Temporal and Spatial Patterns Peter J. Tango<sup>2</sup> Rebecca R. Murphy<sup>1</sup> Richard Tian<sup>1</sup> Jennifer Keisman<sup>4</sup>



### Closing thoughts: Top 10 qualities for good indicators:

### **Important Indicator Qualities**

- 1. Simple and easy to understand
- 2. Be scientifically well-founded
- 3. Have a reference or threshold value of significance
- 4. Be responsive to changes in the environment
- 5. Show trends over time
- 6. Feasible to measure and report (reasonable cost/benefit ratio)
- 7. Updated regularly with reliable procedures (timely with support of a monitoring program)
- 8. Adequately documented, known quality
- 9. Be useable by the community
- 10. Policy relevant


# Additional indicator qualities to potentially keep in mind

- Clear in value: no uncertainty about which direction is good and which is bad.
- Clear in content: easily understandable, with units that make sense.
- Compelling: interesting, exciting, suggestive of effective action.
- Policy relevant: for all stakeholders in the system.
- Feasible: measurable at reasonable cost.
- Sufficient: not too much or too little information.
- Timely: compliable without long delays.
- Appropriate in scale: neither over- nor under-aggregated.
- **Democratically** chosen and accessible.
- Supplementary: include what people cannot measure for themselves.
- **Participatory:** make use of what people can measure for themselves and compile it to provide geographic or time overviews.





### Issue

- The public frequently looks to the Chesapeake Bay Program Office of Communications for updates about The Dead Zone.
- There is no outcome in the Watershed Agreement specifically calling out The Dead Zone event, and no single group tasked with specifically tracking and reporting on the Dead Zone phenomenon.
  - Therefore, collaborative discussions happen but ad hoc annual reporting occurs.
- Requesting input, guidance, discussion on
  - A need to organize a communications strategy within the CBP regarding this annual Bay event
  - Directing any new analyses and resources to produce independent CBP products for reporting, or
  - We are fine, continue pulling together stories sometime after the Dead Zone goes away each year




# Bay Health Status: Dissolved Oxygen Time Series Water Quality Standards Criteria Attainment for All Tidal Waters 1985-2012





#### The Chesapeake Bay's Shrinking 'Dead Zones'

As nutrient pollution into the bay decreases, fewer algae blooms mean there is more dissolved oxygen available in the water available for fish and other marine animals that require oxygen to survive.



Source: NPR analysis of Chesapeake Bay Program and USGS data; Rebecca Murphy/Chesapeake Bay Program

Credit: Leanne Abraham/NPR





ABOUT EDUCATION RESEARCH CAMPUSES APPLYING SCIENCE DIRECTORY NEWS & EVENTS CONTACT

## LARGE 'DEAD ZONE' FORECAST FOR CHESAPEAKE BAY AFTER WET WINTER AND SPRING



June 12, 2019

**Chesapeake Biological Laboratory** 

Ecologists from the University of Maryland Center for Environmental Science and the University of Michigan are forecasting a large Chesapeake Bay "dead zone" in 2019 due to well-above-average river flows associated with increased rainfall in the watershed since last fall.



#### News

- > Boating
- > Fishing
- > Forestry
- > Hunting
- > Lands
- > Parks
- > Police
- > Waters
- > Wildlife

#### **Media Tools**

- > Contact Us
- > News
- > Newsletter
- Magazine
- > Mobile App
- Online Store

#### July 2019 Hypoxia Report

August 2, 2019


#### Maryland Department of Natural Resources

monitoring data show that dissolved oxygen conditions in the Maryland portion of the Chesapeake Bay mainstem were larger than average in July, as the result of many factors. The hypoxic water volume (areas with less than 2 mg/l oxygen) was 1.92 cubic miles in early July and 2.01 cubic miles in late July. The early July 1985-2018 average hypoxic volume is 1.36 cubic miles, and late July is 1.34 cubic miles. The most recent hypoxic volumes were respectively the fourth-largest



for early July and second-largest for late July since 1985.

Larger than average hypoxia was predicted for this summer in part due to massive rainfall causing high flows into the Chesapeake Bay the past year, which delivered higher nutrient loads. Those nutrients fuel algal blooms, which die and are consumed by bacteria, which then deplete oxygen in bottom waters. Also, average winds were generally low in July which prevented oxygen from mixing into deeper waters.



https://news.maryland.gov/dnr/2019/08/02/july-2019-hypoxia-report/


























Chesapeake Bay Program Dead Zone reporting is commonly a synthesis of agency and institution reports and research...




### "Hypoxic Volume" – what does that mean?




# Bay Health Status: Dissolved Oxygen Time Series Water Quality Standards Criteria Attainment for All Tidal Waters 1985-2012




"Hypoxic Volume" – what does that mean?
In this case it means July average volume of water <2mg/L for the mainstem Bay




# Indicators: What the Public Sees of Our Water Quality Monitoring Program





Chesapeake Bay Hypoxia Summer 2012




Is it satisfactory to the CBP to continue Dead Zone reporting in an ad hoc messaging fashion?

Is there a need to establish a separate analysis prepared by CBP analysts to identify as "The Chesapeake Bay Program Dead Zone report?"

Is there a need to organize an annual collaborative newsletter at a set time each year integrating forecasts-summer conditions-final findings?

#### Potomac Estuary Wastewater Treatment and Summer Surface DO Concentrations at Woodrow Wilson Bridge Station





CHESAPEAKE BAY DAILY

Background

Contact Information

Hypoxia (Dissolved Oxygen)

Dead Zone Size

Depth to Low Oxygen

Hypoxia Line Plots

Bay-wide Salinity and Temperature

Focused Salinity and Temperature Forecasts

Chlorophyll

Striped Bass Habitat

Ocean Acidification

Home > Research & Services > Topics > Dead Zones > Forecasts > Chesopeake Bay Daily > hygoxic\_volume

#### Volume of Hypoxic Water


#### Chesapeake Bay


▶ Quick Summary

Year

#### 2019 Dead Zone Size

The amount of hypoxia this summer was predicted to be quite large, as a result of high river inflows to the Bay this year. The amount of hypoxia in the Bay is expected to increase from spring into summer and then decrease as summer progresses into fall. However, short-term weather will affect the amount of hypoxia in the Bay. Check back to see how the size of the dead zone increases seasonally and how daily weather changes the amount of hypoxia. Notable weather that may impact the amount of hypoxia are very windy days or periods of very calm wind. The image below will be continually updated throughout 2019 based on the daily forecast model.






Hypoxic Volume (HV) Metrics for Recent Years for Comparison to 2019 Forecast

 Maximum Daily HV
 Average Summer HV
 Hypoxic Duration
 Total Annual HV


 [km³]
 [km³]
 [days]
 [km³ days]




Home / Facts / What is a dead zone?

### What is a dead zone?

"Dead zone" is a more common term for hypoxia, which refers to a reduced level of oxygen in the water.




Hypoxic zones are areas in the ocean where the oxygen concentration is so low that animals can suffocate and die, and as a result are often called "dead zones." The largest hypoxic zone in the United States, and the second largest hypoxic zone worldwide, forms in the northern Gulf



# Bay Health Status: Dissolved Oxygen Time Series Water Quality Standards Criteria Attainment for All Tidal Waters 1985-2012

This is roughly what we show the public on Chesapeake Progress and how we describe it




## Bay Health Status: Dissolved Oxygen Time Series Water Quality Standards Criteria Attainment for All Tidal Waters 1985-2012

This is roughly what we show the public on Chesapeake Progress

and how we describe it









# Make the issue of low oxygen relatable to the public?

### Call it a **Dead Zone**



MODELING PUBLICATION FEDERAL


#### 2022 Chesapeake Bay Dead Zone Report Released

NOVEMBER 16, 2022 - ANCHOR QEA AND THE VIRGINIA INSTITUTE OF MARINE SCIENCE


Chesapeake Bay, the largest estuary in the United States and the third largest in the world, is particularly vulnerable to human impacts. In recent decades, low dissolved oxygen and changing acidification have severely affected the Bay's ecosystem. A dead zone—an area of water with little to no oxygen—can form when excess nutrients, such as nitrogen and phosphorous from fertilizer, wastewater, and urban runoff, enter the water. Scientists refer to these oxygen-poor waters as "hypoxic," meaning the oxygen concentration is less than 2 milligrams per liter, which is too low for aquatic life to thrive.

In collaboration with the Virginia Institute of Marine Science (VIMS), Anchor QEA operates the Chesapeake Bay Environmental Forecast System (CBEFS) computer model, which both predicts dissolved oxygen concentrations and forecasts environmental conditions in the Bay every day. Anchor QEA and VIMS recently released a 2022 Chesapeake Bay Dead Zone Report that summarizes the severity of the dead zone during the summer of 2022 and compares these findings to data collected from 1985 to 2021, with a focus on the previous 5 years.





# Capture public attention:





Killer bees are spreading across the US – and are being drawn to northern states because of warming temps

Story by Erin Keller • 5h • ⑦ 2 min read Summarize

Africanized honey bees, infamously known as "killer bees," are rapidly spreading across the United States, having

already been confirmed in 13 states from California and Arizona to Florida, Georgia, and as far east as Alabama.

