

**OLD DOMINION UNIVERSITY**

<sup>1</sup>Department of Biological Sciences  
Old Dominion University, Norfolk, Virginia 23529

<sup>2</sup>Versar, Incorporated  
9200 Rumsey Road, Columbia, Maryland 21045

<sup>3</sup>Virginia Institute of Marine Science  
College of William and Mary, Gloucester Pt., Virginia, 23062

**Preliminary evaluations of secondary productivity estimates as indicators of the ecological value of the benthos to higher trophic levels in Chesapeake Bay**

Prepared by

Principal Investigators:

Daniel M. Dauer<sup>1</sup>  
Michael F. Lane<sup>1</sup>  
Roberto J. Llansó<sup>2</sup>  
Robert Diaz<sup>3</sup>

Submitted to:

Virginia Department of Environmental Quality  
629 East Main Street  
Richmond, Virginia 23230

December, 2011

## CONTENTS

INTRODUCTION .....	2
METHODS.....	3
Probability-based monitoring .....	3
Benthic Secondary Production.....	4
RESULTS AND DISCUSSION .....	5
Comparison of Edgar's versus Brey's estimates of benthic secondary productivity.....	5
Benthic productivity - patterns and relationships with physico-chemical and benthic metrics ..	5
<i>B-IBI Habitat Types</i> .....	5
<i>CBP Benthic Monitoring Strata</i> .....	6
<i>CBP monitoring Segments</i> .....	6
<i>CBP monitoring Segments - Low DO and Contaminant Effects</i> .....	7
Low Dissolved Oxygen and CBP Segment Benthic Productivity .....	7
Sediment Contaminants and CBP Segment Benthic Productivity .....	8
SUMMARY .....	9
LITERATURE CITED .....	10

## INTRODUCTION

Benthic macrofauna are important components of estuaries because they affect the physical and chemical properties of the sediment and water column, influence nutrient cycling, are food sources for higher trophic levels, are themselves commercially important and because their sedentary nature and life history characteristics expose them directly to both natural and man-made disturbances making them good indicators of those effects on estuarine systems (Reish, 1973; Boesch, 1973; Boesch, 1977; Pearson and Rosenberg, 1978; Bilyard, 1987; Gray, et al., 1988; Dauer, 1993). As a result, a common approach to environmental assessment in estuarine systems is to develop indicators of macrobenthic community health that are measures of diversity, abundance, biomass and/or metrics related to sensitive or tolerant macrobenthic species (see review by Borja and Dauer, 2008). Such assessments, however, rarely involve evaluations of benthic secondary production (Dolbeth et al., 2010).

Macrobenthic secondary production can be defined as the amount of organic matter incorporated into macrobenthic (>0.5mm) invertebrates per unit time and typically expressed per unit area. As such, these estimates would represent the amount of energy processed by these organisms from organic detritus and other sources and made available to higher trophic levels such as, crabs, bottom-feeding fish, diving ducks and shallow-water foraging birds. With respect to environmental management, measures of secondary production can be used to assess (1) the value of benthos as a food source for higher trophic levels (Asmus, 1987; Franz and Tanacredi, 1992; Tumbiolo and Downing, 1994; Wilson, 2002; Cusson and Bourget, 2005; Dolbeth et al., 2010); (2) the effects of anthropogenic inputs such loads of nutrients, sediment and contaminants on benthic communities (Steimle, 1985; Dolbeth 2003; Dolbeth 2007; Dauer et al. 2000; Bolam et al., 2011); (3) the efficacy of restoration and environmental management (Borja et al. 2008, 2010, 2011) particularly in an adaptive management and adaptive monitoring sense (Borja and Dauer, 2008); and (4) the impacts of climate change on macrobenthic communities (Dolbeth et al., 2007).

The primary objective of this project was to develop an analytical tool that management could use to assess the ecological value of the benthos for higher trophic levels. Ideally any tool developed should be simple to implement, adaptable to both fixed and probability monitoring programs and applicable across multiple spatial and temporal scales. Direct evaluations of secondary productivity involve time-consuming cohort or size-based methods that require multiple measurements of the populations in question over time (Dolbeth et al., 2005; Cusson and Bourget, 2005). However, empirical regression models have been developed that provide reasonable alternatives to these techniques when direct measurements are not an option (e.g. Schwinghamer et al., 1986; Edgar, 1990; Tumbiolo and Downing, 1994; Brey, 1999; Brey, 2001). These relationships incorporate benthic population metrics biomass, abundance, mean individual weights, etc.), environmental metrics (temperature, water depth), and taxon-specific metrics to adjust for body-size dependent turn-over rates to estimate production rates (or P/B ratios can be used to obtain production rates). Of the available empirical relationships we evaluated two approaches to calculate benthic secondary production in Chesapeake Bay – those of Edgar (1990) and Brey (2001) described in detail in the Methods section. Edgar's

(1990) equation was selected because it has been used in several other studies conducted in Chesapeake Bay (Diaz and Schaffner 1990) and elsewhere (Wilbur and Clarke, 1998; Gillet, 2011; Kersey, 2011). Brey's (2001) equation was selected because it incorporates variation related not only to temperature but also to sample depth and to effects relating to taxonomic composition that were not incorporated into Edgar's model and because comparisons of different empirical models found that Brey's model generally performed better than others (Dolbeth et al., 2005; Cusson and Bourget, 2005; Bolam et al., 2011). Finally, both of these equations can be easily calculated using data directly available from existing Chesapeake Bay Program data sets.

This study presents our preliminary evaluations of both of these allometric equations for use in developing an assessment tool using benthic secondary production in Chesapeake Bay. We present estimates of average rates of secondary production at three spatial scales for the entire tidal portion of the Chesapeake Bay watershed as follows: (1) the 7 different B-IBI Weisberg habitat types (see Weisberg, et al., 1997); (2) the ten benthic sampling strata of the stratified-random benthic of the monitoring program in Chesapeake Bay (see Dauer and Llansó 2003) - plus an eleventh stratum from aggregated sites collected in the Elizabeth River watershed; and (3) the 73 Chesapeake Bay Program segments used a management tool to report overall condition in specific regions (*i.e.* 305(b)/303(d) reporting) (see, Llansó et al., 2009). Results from both equations were compared to values in the literature from Chesapeake Bay and similar estuaries.

Our intent is to develop a benthic secondary productivity assessment tool to help in assessing ecological status patterns using the Benthic Index of Biotic I (B-IBI) (Weisberg et al., 1997). The B-IBI has been used effectively in Chesapeake Bay to: (1) identify impaired waters for Federally mandated management reports (Llansó et al., 2009); (2) estimate the extent of areal degradation of the watershed (Dauer and Llansó, 2003); and (3) relate benthic community condition to water quality, sediment quality and watershed stressors on a bay-wide scale (Dauer et al., 2000). We believe that complimentary studies investigating benthic secondary production will enhance adaptive management strategies at multiple spatial scales both for macrobenthic communities and higher trophic levels.

## METHODS

### Probability-based monitoring

This study presents samples collected as part of the Chesapeake Bay Program's probability-based benthic monitoring program from 1996 through 2009 coupled with probability-based samples collected as part of Virginia DEQ's Elizabeth River Monitoring Program conducted from 1999 through 2007. The Chesapeake Bay Program allocated samples by assigning 25 random locations on an annual basis into 10 separate strata: (1) three in the Chesapeake Bay Mainstem; (2) five in major tributaries to the Mainstem; (3) two in Maryland eastern and western shore tributaries (see Dauer and Llansó 2003). For the Elizabeth River Monitoring Program, the main

sampling strata corresponded to main subwatersheds of the Elizabeth River i.e. Western, Southern, and Eastern branches, the Lafayette River, the Elizabeth River Mainstem, as well as, several small contaminated tidal creeks sampled as additional strata during 1999 and 2002 (Dauer 2000, 2009). As with the CBP 25 random locations were assigned on an annual basis to each stratum.

All samples were collected during the Benthic IBI index period (Weisberg et al., 1997) using a Young grab with a surface area of 440 cm<sup>2</sup>. Each sample was sieved on a 0.5 mm screen and preserved in the field. Samples were sorted, enumerated and identified to the lowest possible taxon. Ash-free dry weight biomass was determined for each taxon.

### Benthic Secondary Production

We assessed the applicability of two general allometric equations as indicators of benthic secondary production. The first was that of Edgar (1990) which relates per sample daily macrobenthic production  $P$  to per sample standing crop biomass  $B$  in mg ash free dry weight (AFDW) and water temperature  $T$  in °C as follows:

$$P = 0.0049 \times B^{0.80} T^{0.89}$$

Daily per sample estimates of production in g C were estimated for each individual taxon in the community and then summed across all taxa to obtain a daily total community production rate. Daily production rates were converted to annual rates per m<sup>2</sup> by first multiplying the daily rates by 365 d/yr and dividing the result by the total area of the sample in m<sup>2</sup> to obtain g C/m<sup>2</sup>/yr.

The second equation used was a modified version of Brey's (2001) formula for estimating P/B ratios based on mean body mass per individual  $w$  expressed in  $\mu$ g, sample depth  $D$  in meters, temperature in °K and several discrete (dummy) variables which took the following form:

$$\log_{10}(P/B) = 7.947 - 2.294 \times \log_{10}(w) - (2409.856 \times 1/T) + (0.168 \times 1/D) + (0.194 \times Subtid) + (0.180 \times InfEpi) + (0.174 \times Tax1) - (0.188 \times Tax2) + (0.330 \times Tax3) + (582.851 \times \log_{10}(w) \times 1/T)$$

*Subtid* is a dummy variable that increases the P/B ratio if the organism is found in a subtidal habitat (i.e. a depth of > 1 meter) while *InfEpi* is set to 1 if the organism is infaunal also resulting in an increase in the P/B ratio. *Tax1*, *Tax2* and *Tax3* are dummy variables that identify specific effects on P/B ratio associated with membership in different taxonomic groups and that are set to 1 if the organism is: (1) an annelid or crustacean; (2) an echinoderm or (3) an insect, respectively, and 0 if otherwise. These terms result in an increase in P/B ratio for annelids, crustacean and insect species and a decrease in P/B ratio for echinoderm species. The original version of the formula includes additional dummy variable terms that do not apply to the samples collected in this study (e.g. regarding epifaunal mobility and lake habitat effects).

P/B ratios for individual species were estimated by first dividing the AFDW per sample by the total number of individuals to obtain an average mass per individual in g C. These values were then converted to the required units (kJ) using taxonomic group specific conversion factors provided by Brey (2001) (see Appendix A). The species level mass values in combination with depth and temperature recorded at the time of collection were used to calculate  $\log_{10}$  transformed P/B ratios using the equation. By converting the ratio back to a linear scale (i.e. raising as an exponent to 10), the linear P/B ratios obtained were then multiplied to the mean standing crop biomass (per  $\text{m}^2$ ) to obtain an estimate of production per unit area and time for each species. We assumed that the standing crop values per sample were representative of one year of benthic community biomass and that the resultant productivity estimates were in units of  $\text{g C}/\text{m}^2/\text{yr}$  for each species. Total community production for a given site was the summation of all taxa specific production values.

## RESULTS AND DISCUSSION

### Comparison of Edgar versus Brey estimates of benthic secondary productivity

Edgar secondary productivity estimates were generally higher than those of the Brey estimates (Figure 1). The Edgar estimates of productivity were linearly related to biomass with an  $R^2$  value of 0.96 (Figure 2). Brey estimates were poorly related to biomass in a linear manner with an  $R^2 = 0.54$  (Figure 3). Benthic biomass values in Chesapeake Bay are not typically greater than  $100 \text{ gC}/\text{m}^2$ . Biomass values  $> 100 \text{ gC}/\text{m}^2$  (Figure 4A) again show Edgar values for secondary productivity much higher the Brey values with the same pattern for biomass values  $> 5 \text{ gC}/\text{m}^2$  that account for the vast majority of biomass values in Chesapeake.

In essence Edgar values for benthic secondary productivity are basically a simple linear transformation of biomass values. For macrobenthic communities that contain species (1) with a diversity of life spans, (2) very different allocations of energy into protective coverings or other refugia from predation (e.g. high versus low mobility), and (3) taxon-specific differences in P/B ratios that are best explained by natural history traits (Cusson and Bourget, 2005), the linear relationship to biomass of the Edgar values seems ecologically unreasonable.

In addition the Brey equation contains many more variables than Edgar's equation that are likely to affect secondary productivity and Brey values are more consistent with the review of production marine benthic habitats of Cusson and Bourget (2005). Finally our results Brey values were also consistent with estimates provided by Diaz and Schaffner (1990) for various benthic habitat types in Chesapeake Bay (see Appendix B). As such, further results of secondary production presented will be from Brey's equation.

## Benthic productivity - patterns and relationships with physico-chemical and benthic metrics

### ***B-IBI Habitat Types***

In developing the benthic index of biotic integrity, Weisberg et al. (1997) identified seven benthic habitat types in Chesapeake Bay – tidal freshwater (TF), oligohaline (OH), low mesohaline (LMH), high mesohaline mud (HMHM), high mesohaline sand (HMHS), polyhaline mud (PHM) and polyhaline sand (PHS). Benthic community metrics were selected and scored for each of these seven habitat types. For the 3,918 samples used in our study Table 1 summarizes the physico-chemical parameters and Table 2 presents Brey's secondary productivity estimates as well as community level biomass, abundance, B-IBI values and species per sample. In Figures 5-9 the data for these benthic biotic parameters are presented as well as the Shannon Index of informational diversity in Figure 10.

Both benthic secondary productivity and biomass were higher in the three lowest salinity habitat types (TF, OH, LMH)(Figures 5 and 6). This pattern is not obvious from Figure 3 where the highest benthic secondary productivity, e.g.  $> 200 \text{ gC/m}^2/\text{yr}$ , occurs with biomass from  $10 - 700 \text{ gC/m}^2$ . These three lowest salinity habitats had the lowest species richness (Table 2 and Figure 9) and Shannon diversity (Figure 10) as expected from the general Remane Curve relationship for estuarine-transition waters but also lower abundance than the three highest salinity habitat types (HMHS, PHM, PHS) (Figure 7).

### ***CBP Benthic Monitoring Strata***

Since 1996 the Chesapeake Bay Benthic Monitoring Program has presented data on benthic community condition by randomly sampling all tidal waters of the Bay. Allocation of samples is random-stratified approach with each of ten strata allocated 25 random samples (sites) each year (see Figure 11 for the ten strata). The 3,918 samples used in our study were summarized by the stratum of collection. An eleventh stratum for the Elizabeth River watershed was added to the analyses in this section (ELR). The physico-chemical parameters are summarized in Table 3 and the biotic parameters in Table 4. In Figures 12-16 the data for these benthic biotic parameters are presented as well as the Shannon Index of informational diversity in Figure 17.

Benthic secondary productivity varied widely with highest levels in upper mainstem of the Bay (UPB) and the lowest in the Patuxent River (PXR) (Figure 12). As such the benthic secondary productivity and biomass are plotted on a semilog scale in Figures 18 and 19, respectively. The semilog plots emphasize the disconnect in several strata between estimates of benthic secondary productivity and biomass. For example the James River had a much higher biomass than the Elizabeth River ( $4.81$  versus  $1.61 \text{ gC/m}^2$ ) yet the Elizabeth River secondary productivity is comparable to that of the James River ( $18.09 \text{ gC/m}^2/\text{yr}$  in the James and  $18.65 \text{ gC/m}^2/\text{yr}$  in the Elizabeth River).

### ***CBP monitoring Segments***

The Chesapeake Bay Program (CBP) divides the Bay into 73 segments (Figure 20). The 3,918 samples used in our study were summarized by CBP segments. The physico-chemical parameters are summarized in Table 5 and the biotic parameters in Table 6.

The Brey's benthic secondary productivity values for each CBP segment are summarized in Figures 21 and 22. The highest levels of benthic secondary productivity are all in Maryland and typically in TF and OH segments that have significant populations of the bivalve species *Corbicula fluminea*, *Rangia cuneata* or *Gemma gemma* (ELKOH, GUNOH, NANOH, POTTF, POTOH, MIDOH, CHSOH CB1TF, CB2OH and CB4MH).

The second most productive segments were (1) the TF sections of the Virginia tributaries (JMSTF, PMKTF, MPNTF, RPPTF), (2) several OH sections, primarily in smaller tributaries and in Maryland (CHKOH, JMSOH, PAXOH, POCOH, SASOH, BSHOH), (3) several mesohaline small Maryland tributaries (WSTMH, LCHMH, EASMH), (4) the mouth of the Bay (CBPH8) and (5) two segments in the heavily impacted Elizabeth River watershed (SBEMH and EBEMH).

The third group of segments had secondary productivity values between 10 to <20 gC/m<sup>2</sup>/yr. This group included (1) three sections of the Mainstem of the Bay bordered upstream (CB3MH) and downstream (CB6PH, CB7PH) of the Maintsem section (CB5MH) typically subjected to bottom low dissolved oxygen events, (2) all of the Rappahannock River mainstem downstream of the TF section (RPPOH, RPPMH), (3) all of the York River downstream of the TF sections, including MobJack Bay (MPNOH; PMKOH, YRKM, YRKP, MOBPH), (4) all of the James River mainstem downstream of the TF and OH segments (JMSMH, JMSPH), (5) most of the Choptank River (CHOMH2, CHOOH, CHOTF), (6) several mesohaline small Maryland tributaries (MAGMH, PATMH, SOUMH, CHSMH, WICMH, RHDH), and (7) three segments in the heavily impacted Elizabeth River watershed (ELIPH, WBEMH, LAFMH).

The final group of segments had secondary productivity values <10 gC/m<sup>2</sup>/yr and included (1) a single segment in the Mainstem (CB5MH), (2) numerous mesohaline segments in Maryland (POTMH, MANMH, SEVMH, BIGMH, FSBMH, HNGMH, TANMH, POCMH, PAXMH, NANMH, CHOMH1), (3) two mesohaline segments in Virginia (CRRMH, PIAMH), and (4) two lower salinity segments in Maryland (PAXTF, BACOH).

### ***CBP monitoring Segments - Low DO and Contaminant Effects***

The major stressors of the macrobenthic communities of Chesapeake Bay are (1) bottom low dissolved oxygen (driven primarily by excess primary production and a resulting imbalance in aerobic versus anaerobic metabolism at the ecosystem level) and (2) sediment contamination (modified by abiotic chemical and biochemical amelioration of toxic effects, as well as bioturbation effects upon bioavailability). However, in examining the patterns of benthic secondary productivity within the CBP segments two concerns were obvious – the unexpectedly high benthic secondary productivity (1) in the Mainstem segment CB4MH



subjected to periodic and acute low dissolved oxygen events and (2) within the CBP segments in the Elizabeth River watershed (SBEMH, EBEMH) subjected to long-term, chronic exposure to sediment contaminants, primarily high levels of PAHs (Dauer and Llanso 2003).

#### Low Dissolved Oxygen and CBP Segment Benthic Productivity

The effects on the benthos of low oxygen events are globally widespread (Diaz and Rosenberg 1995) and well documented in Chesapeake Bay (Pihl et al., 1991; Dauer et al. 1992, 1993; Dauer 1993). Marine and coastal ecosystems with hypoxic and or anoxic bottom water conditions have low annual secondary production (Diaz and Rosenberg 2008). Diaz and Rosenberg (2008) estimated for Chesapeake Bay that ~10,000 MT C is lost because of hypoxia each year, representing ~5% of the Bay's total secondary production. Low dissolved oxygen events alter benthic energy flow pathways with more energy diverted into microbial pathways including in essential benthic habitats that are nursery and recruitment areas.

CB4MH has spatially extensive low dissolved during the summer especially in the channel region (Figure 23); however, higher dissolved oxygen levels supportive of benthos are found on the shallower shoals, particularly on the eastern side of the segment. Closer examination within CB4MH indicates that the high segment level secondary productivity values is driven by shallow water samples (<8m) highly dominated by the small but productive bivalve *Gemma gemma*. Figures 24 and 25 show the distribution of secondary productivity in CB4MH with various depth intervals.

Lower rates of benthic production associated with lower levels of dissolved oxygen were clearly shown in other segments, for example, the lower Potomac River (POTMH). In this segment, very low bottom dissolved oxygen levels occur each summer especially at great water depths. Benthic secondary productivity was by far the highest in depths shallower than 5 m (Figure 26). Benthic secondary productivity declined from levels of 243 gC/m<sup>2</sup>/yr in water depths <5m to <5 gC/m<sup>2</sup>/yr in water depths >20m. The percentage of azoic samples (no benthos found) increased with depth with over 70% of the benthic sample in >20m being azoic.

In general CBP segments with lower dissolved oxygen levels has lower benthic secondary productivity. For example, there were 12 CBP segments (excluding CBP segments with < 10 samples) with a summer mean bottom dissolved oxygen value > 5mg/l (see values in Table 5). (CBP segments with > 10 samples were excluded). These segments were (1) Mainstem segments CB3MH, CB5MH and CB6PH, (2) the lower reaches of the Potomac River (POTMH) and the York River (YRKPH), (3) the oligohaline segments of both source tributaries of the York River (MPNOH, PMKOH), and (4) several mesohaline segments in highly urbanized regions of Maryland - the Patapsco River (PATMH), the Magothy River (MAGMH), and the Severn River (SEVMH) – and Virginia's Elizabeth River – the Southern Branch (SBEMH), and the Eastern Branch (EBEMH). Ten of the twelve segments were in the two lowest ranges of productivity (see Figure 27), except for the two segments in the Elizabeth River (the Southern Branch - SBEMH, and the Eastern Branch - EBEMH).

### Sediment Contaminants and CBP Segment Benthic Productivity

Two of the segments in the Elizabeth River were in the second highest benthic productivity level (Figure 21) although at the lower end of the scale (Figure 22). Generally sediment contaminant levels are closely related to near-field levels of urban land-use. As such Figure 22 is modified in Figures 28 and 29 to indicate the five segments of the Elizabeth River watershed (ELIPH, LAFMH, WBEMH, EBEMH, SBEMH), and the ten segments of the Maryland Western Tributaries (BSHOH, GUNOH, MIDOH, BACOH, PATMH, MAGMH, SEVMH, SOUMH, RHDMH, WSTMH) all of which are relatively small segments by benthic surface area and have shorelines with high levels of urban land-use. These small urbanized segments have values of benthic secondary productivity in all four intervals shown in Figure 21.

In order to better understand the dynamics of benthic secondary production and the structural benthic metrics, four CBP segments were further compared - JMSMH, POTMH, PATMH and SBEMH. All are mesohaline regions with the James River segment (JMSMH) having no bottom low dissolved oxygen events and no known sediment contaminant levels of concern. The Potomac River segment (POTMH) has significant low dissolved oxygen events during the summer but no known sediment contaminant levels of concern. Both the Patapsco River segment (PATMH) and the Elizabeth River segment (SBEMH) are characterized by levels of sediment contaminants that have significant biological effects. The highest benthic secondary productivity (Figure 30) was in the two contaminant influenced segments (PATMH, SBEMH) a pattern not reflective of the biomass pattern (Figure 31). The lowest benthic secondary productivity value was the Potomac River which also had relatively high and variable biomass value especially compared to the James River segment (JMSMH). The two sediment contaminated segments had similar levels of benthic secondary productivity (Figure 30) but very different biomass values (Figure 31). The high biomass values in the Patapsco River segment are driven by bivalve species *Macoma balthica* and *Rangia cuneata* that are rare in the Southern Branch of the Elizabeth River. The high levels of benthic secondary productivity in the Southern Branch is driven by the very high abundances (Figure 32) and high productivity of the polychaetes *Mediomastus ambiseta* and *Streblospio benedicti* (Tables 7- 9). B-IBI values (Figure 33), Species richness (Figure 34), and the Shannon diversity index (Figure 35) follow the expected pattern with (1) the lowest values of these three metrics in the Potomac River segment where numerous azoic samples were collected (Figure 26), (2) highest values in general at the James River segment, and (3) intermediate at the two sediment contaminated segments in the Patapsco and Elizabeth rivers.

In assessing species-level influences upon benthic secondary productivity, the comparison between JMSMH and SBEMH is indicative of the disconnect between benthic community abundance (Figure 32), biomass (Figure 31) and benthic secondary productivity (Figure 30). Abundance of both segments was dominated by annelids with the top ten density dominant being annelid species in SBEMH and seven of the top ten annelids in JMSMH (Table 8). However abundances of the polychaete species *Mediomastus ambiseta* and *Streblospio benedicti* were much higher in SBEMH. Biomass dominants in SBEMH were also primarily annelids (nine of ten); however, in JMSMH the top biomass dominants included four bivalve

species (*Mercenaria mercenaria*, *Macoma balthica*, *Macoma mitchelli*, *Rangia cuneata*), one gastropod species (*Polinices duplicatus*) and one amphipod species (*Leptocheirus plumulosus*) (Table 9). Finally when benthic secondary productivity rates were calculated the primary differences were (1) the great importance in both segments of productivity rates of polychaete species (*Mediomastus ambiseta*, *Streblospio benedicti*, *Leitoscoloplos* spp., *Heteromastus filiformis*, *Paraprionospio pinnata*, *Neanthes succinea*), (2) the great productivity rates in JMSMH of bivalve species *Macoma balthica* and the amphipod *Leptocheirus plumulosus* that are rarely collected in SBEMH, and (3) the higher productivity rate in SBEMH of the polychaete species *Laonereis culveri* that is rarely collected in JMSMH.

### **Summary**

The relationship between benthic community condition as measured by the B-IBI and rates of benthic secondary productivity is not simple (Figure 36). High secondary productivity can be associated with either low BIBI levels characterized as severely degraded ( $BIBI \leq 2.0$ ) or with BIBI values considered undegraded ( $BIBI \geq 3.0$ ). In addition, regions with high levels of sediment contaminants such as the Patapsco River and Elizabeth River can have unexpectedly high levels of benthic secondary productivity

The next steps in using benthic secondary productivity estimates is to develop a protocol to reflect the actual availability of the benthic production to higher trophic levels. Important ecological factors are (1) protective coverings such as molluscan shells and crustacean exoskeletons that reduce predation, (2) depth of dwelling within the sediment that might provide a refuge from predation, (3) body size factors that affect strength of protective coverings and/or age-related sediment depth dwelling location, and (4) general behaviors that can modify susceptibility to predation, e.g. rapid motility.

In natural ecosystems, local species diversity and productivity are regulated by a myriad of interacting factors at a variety of temporal and spatial scales. Ecological status assessment is essential to direct maintenance, protection, and/or restoration efforts in regard to marine and estuarine ecosystem services. Parameters that assess ecosystem structure are widely used, diverse, and often profligate. Parameters that assess ecosystem function should be less variable and better understood in directing ecosystem management decisions. Thus assessment of benthic secondary productivity estimates as a management assessment tool deserves serious further consideration, development as a tool and application in management decisions.

## Literature Cited

- Alden, R.W. III, S.B. Weisberg, J.A. Ranasinghe, and D.M. Dauer. 1997. Optimizing temporal sampling strategies for benthic environmental monitoring programs. *Marine Pollution Bulletin* 34:913-922.
- Asmus, H., 1987. Secondary production of an intertidal mussel bed community related to its storage and turnover compartments. *Marine Ecology Progress Series*. 39:251-266.
- Bilyard, G.R., 1987. The value of benthic infauna in marine pollution monitoring studies. *Marine Pollution Bulletin*. 18:581-585.
- Boesch, D.F., 1973. Classification and community structure of the macrobenthos in the Hampton Roads area, Virginia. *Marine Biology*. 21:226-224.
- Boesch, D.F. 1977. A new look at zonation of the benthos along the estuarine gradient. pp. 285-308, In *Marine Benthic Dynamic*, K.R. Tenore and B.C. Coull, eds., University of South Carolina Press, Columbia SC.
- Bolam, S.G., J. Barry, T. Bolam, C. Mason, H.S. Rumney, J.E. Thain, and R.J. Law. 2011. Impacts of maintenance dredged material disposal on macrobenthic structure and secondary productivity. *Marine Pollution Bulletin* 62:2230-2245.
- Borja, A. and D.M. Dauer. 2008. Assessing the environmental quality status in estuarine and coastal systems: comparing methodologies and indices. *Ecological Indicators* 8: 331-337.
- Borja, A, S.B. Bricker, D.M. Dauer, N.T. Demetriades, J.G. Ferreira, A. T. Forbes, P. Hutchings, X. Jia, R. Kenchington, J.C. Marques and C. Zhu. 2008. Overview of integrative tools and methods in assessing ecological integrity in estuarine and coastal systems worldwide. *Marine Pollution Bulletin* 56: 1519-1537.
- Borja, A, D.M. Dauer, M. Elliott, and C.A. Simenstad. 2010. Medium and long-term recovery of estuarine and coastal ecosystems: patterns, rates and restoration effectiveness. *Estuaries and Coasts* 33:1249–1260.
- Borja, Á, D. M. Dauer, and A. Grémare. 2011. The importance of setting targets and reference conditions in assessing marine ecosystem quality. *Ecological Indicators*. 12:1-7.
- Brey, T. 1999. A collection of empirical relations for use in ecological modelling. *ICLARM Quarterly* 22:24-28.
- Brey, T. 2001. Population dynamics in benthic invertebrates. A virtual handbook. Version 01.2. <http://www.thomas-brey.de/science/virtualhandbook>.

Cusson, M. and E. Bourget, 2005. Global patterns of macroinvertebrate production in marine benthic habitats. *Marine Ecology Progress Series*. 279:1-14.

Dauer, D.M., 1993. Biological criteria, environmental health and estuarine macrobenthic community structure. *Marine Pollution Bulletin* 26: 249-257.

Dauer, D.M. 1997. Dynamics of an estuarine ecosystem: Long-term trends in the macrobenthic communities of the Chesapeake Bay, USA (1985-1993). *Oceanologica Acta* 20:291-298.

Dauer, D.M. 2000. Benthic Biological Monitoring Program of the Elizabeth River Watershed (1999). Final Report to the Virginia Department of Environmental Quality, Chesapeake Bay Program, 73 pp.

Dauer, D.M. 2009. Benthic Biological Monitoring Program of the Elizabeth River Watershed (2008). Final Report to the Virginia Department of Environmental Quality, Chesapeake Bay Program, 119 pp.

Dauer, D.M. and R.W. Alden III. 1995. Long-term trends in the macrobenthic communities of the lower Chesapeake Bay (1985-1991). *Marine Pollution Bulletin* 30:840-850.

Dauer, D.M. and R.J. Llansó. 2003. Spatial scales and probability based sampling in determining levels of benthic community degradation in the Chesapeake Bay. *Environmental Monitoring and Assessment* 81:175-186.

Dauer, D.M., Luckenbach, M.W. and A.J. Rodi, Jr. 1993. Abundance biomass comparison (ABC method): effects of an estuarine gradient, anoxic/hypoxic events and contaminated sediments. *Marine Biology* 116: 507-518.

Dauer, D.M., J. A. Ranasinghe, and S. B. Weisberg. 2000. Relationships between benthic Community condition, water quality, sediment quality, nutrient loads, and land use patterns in Chesapeake Bay. *Estuaries* 23: 80-96.

Dauer, D.M., Rodi, A.J., Ranasinghe, J.A., 1992. Effects of low dissolved oxygen events on the macrobenthos of the lower Chesapeake Bay. *Estuaries* 15, 384– 391.

Diaz, R.J. and R. Rosenberg, R., 1995. Marine benthic hypoxia: a review of its ecological effects and the behavioural responses of benthic macrofauna. *Annual Review of Oceanography and Marine Biology* 33: 245– 303.

Diaz, R.J. and R. Rosenberg, R., 2008. Spreading dead zones and consequences for marine ecosystems. *Science* 32: 926-929

- Diaz, R.J. and L.C. Schaffner. 1990. pp. 25–56, in *Perspectives in the Chesapeake Bay: Advances in Estuarine Sciences*, M. Haire, E. C. Krome, Eds., Chesapeake Research Consortium, Gloucester Point, VA.
- Dolbeth, M., A.I. Lillebø, P.G. Cardoso, S.M. Ferreira, and M.A. Pardal, 2005. Annual production of estuarine fauna in different environmental conditions: An evaluation of the estimation methods. *Journal of Experimental Marine Biology and Ecology* 326:115-127.
- Dolbeth, M., M. A. Pardal, A. I. Lillebø, U. Azeiteiro, and J. C. Marques, 2003. Short- and long-term effects of eutrophication on the secondary production of an intertidal macrobenthic community. *Marine Biology*. 143:1229-1238
- Edgar, G.J., 1990. The use of the size structure of benthic macrofaunal communities to estimate faunal biomass and secondary production. *Journal of Experimental Marine Biology and Ecology* 137:195-214.
- Flint, R.W., 1985. Long-term estuarine variability and associated biological response. *Estuaries* 8:158-189.
- Gillet, D.J. 2010. Effects of habitat quality on secondary production in shallow estuarine waters and the consequences for the benthic-pelagic food web. PhD Dissertation. College of William and Mary, Virginia. 175 p.
- Gray, J. S., M. Ascan, M.R. Carr, K.R. Clarke, R.H. Green, T.H. Pearson, R. Rosenberg, and R.W. Warwick, 1988. Analysis of community attributes of the benthic macrofauna of Frierfjord/Langesundfjord and in a mesocosm experiment. *Marine Ecology Progress Series* 46:151-165.
- Holland, A. E, Mountford, N. K., Hiegel, M. H., Kaumeyer, K. R. and Mihursky, J. A. (1980). Influence of predation on infaunal abundance in upper Chesapeake Bay. *Marine Biology* 57:221-235.
- Llansó, R.J., D.M. Dauer, and J.H. Vølstad. 2009. Assessing ecological integrity for impaired waters decisions in Chesapeake Bay, USA. *Marine Pollution Bulletin* 59: 48-53.
- Llansó, R.J., D.M. Dauer, J.H. Vølstad, and L.S. Scott. 2003. Application of the Benthic Index of Biotic Integrity to environmental monitoring in Chesapeake Bay. *Environmental Monitoring and Assessment* 81:163-174.
- Nixon, S.W. and B.A. Buckley, 2002. "A strikingly rich zone" Nutrient enrichment and secondary production in coastal marine ecosystems. *Estuaries* 25:782-796.
- Pearson, T., and R. Rosenberg, 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanography Marine Biology Annual Review*. 16:229-311.

Pihl, L., S. P. Baden, and R. J. Diaz. 1991. Effects of periodic hypoxia on distribution of demersal fish and crustaceans. *Marine Biology* 108:349-360.

Reisch, D., 1973. The use of benthic animals in monitoring the marine environment. *Journal of Environmental Planning and Pollution Control*. 1:32:38.

Schwinghamer, P., B. Hargrave, D. Peer, C.M. Hawkins, 1986. Partitioning of production and respiration among size groups of organisms in an intertidal benthic community. *Marine Ecology Progress Series* 31:131-142.

Steimle, F.W. Jr., 1985. Biomass and estimated productivity of the benthic macrofauna in the New York Bight: a stressed coastal area. *Estuarine, Coastal and Shelf Science* 21:539-554.

Sturdivant, S.K., 2011. The effects of hypoxia on macrobenthic production and function in the lower Rappahannock River, Chesapeake Bay, USA. PhD. Dissertation. College of William and Mary, Virginia.

Tumbiolo, M.L. and J. Downing, 1994. An empirical model for the prediction of secondary production in marine benthic invertebrate populations. *Marine Ecology Progress Series* 114:165-174.

Wilber, D.H. and D.G. Clarke, 1998. Estimating secondary production and benthic consumption in monitoring studies: a case study of impacts of dredged material disposal in Galveston Bay. *Estuaries*. 21:230-245.

Wilson, J.G., 2002. Productivity, fisheries and aquaculture in temperate estuaries. *Estuarine, Coastal and Shelf Science*. 55:953-967.

Table 1. BIBI habitat types. Physico-chemical parameters. S – salinity, %SC – percent sily-clay, T – bottom water temperature ( $^{\circ}\text{C}$ ), DO – bottom dissolved oxygen (ppm), D – water depth (m). SE – standard error for the previous column. Number of random sites in parentheses.

HABITAT	S	SE	%SC	SE	DO	SE	D	SE	T	SE
Tidal Freshwater (190)	0.2	0.0	60.5	2.5	7.0	0.1	3.7	0.2	24.9	0.2
Oligohaline (353)	2.3	0.1	68.9	1.7	6.9	0.1	3.6	0.1	24.7	0.2
Low Mesohaline (926)	9.3	0.1	60.7	1.1	5.7	0.1	4.4	0.1	25.3	0.1
High Mesohaline Mud (854)	14.9	0.1	82.4	0.5	4.7	0.1	6.8	0.2	25.6	0.1
High Mesohaline Sand (604)	15.0	0.1	9.6	0.4	6.3	0.1	4.3	0.1	25.4	0.1
Polyhaline Mud (518)	21.1	0.1	78.7	0.8	4.7	0.1	7.4	0.2	26.8	0.1
Polyhaline Sand (473)	21.8	0.1	12.3	0.5	5.7	0.1	5.8	0.2	26.3	0.1

Table 2. BIBI habitat types. Biotic parameters. P – Brey's secondary productivity ( $\text{gC}/\text{m}^2/\text{yr}$ ), B – biomass (AFDW  $\text{gC}/\text{m}^2$ ), A – abundance (individuals/ $\text{m}^2$ ), BIBI – benthic index of biotic integrity, S – species per sample. SE – standard error for the previous column. Number of random sites in parentheses.

HABITAT	P	SE	B	SE	A	SE	B_IBI	SE	S	SE
Tidal Freshwater (190)	32.44	4.35	22.0	4.92	3,780	243	3.3	0.08	8.43	0.24
Oligohaline (353)	36.00	3.25	25.9	3.33	3,499	295	2.8	0.04	7.70	0.17
Low Mesohaline (926)	24.55	1.34	12.6	1.29	3,780	295	2.7	0.03	8.37	0.13
High Mesohaline Mud (854)	8.37	0.50	1.4	0.31	2,277	149	2.3	0.03	7.02	0.16
High Mesohaline Sand (604)	16.34	2.09	1.8	0.24	5,920	1,098	2.7	0.03	12.21	0.20
Polyhaline Mud (518)	13.69	0.54	1.2	0.10	4,092	246	2.2	0.03	12.14	0.23
Polyhaline Sand (473)	18.04	0.83	2.9	0.39	4,722	327	3.1	0.04	19.19	0.34



Table 3. CBP benthic Monitoring Program Benthic Strata. Physico-chemical parameters. S – salinity, %SC – percent sily-clay, T – bottom water temperature ( $^{\circ}\text{C}$ ), DO – bottom dissolved oxygen (ppm), D – water depth (m). SE – standard error for the previous column. Number of random sites in parentheses. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.

STRATUM	S	SE	%SC	SE	DO	SE	D	SE	T	SE
UPB	8.10	0.27	63.04	1.79	6.04	0.12	5.76	0.17	24.45	0.09
MWT	8.81	0.22	62.88	1.80	5.51	0.14	4.10	0.17	24.96	0.11
MET	9.82	0.27	51.65	1.88	6.56	0.09	3.63	0.15	24.09	0.11
PMR	10.76	0.31	66.15	1.90	4.82	0.14	7.14	0.23	24.55	0.11
RAP	11.97	0.30	61.68	2.01	5.39	0.10	4.89	0.23	26.21	0.10
PXR	12.01	0.20	61.47	1.86	5.30	0.10	5.64	0.24	25.26	0.09
JAM	12.72	0.44	59.28	1.88	6.05	0.07	4.73	0.21	26.67	0.13
MMS	15.11	0.16	33.08	1.91	6.24	0.12	6.00	0.20	24.66	0.10
YRK	15.47	0.36	59.05	1.86	5.08	0.08	4.79	0.22	27.25	0.08
ELR	20.22	0.17	58.88	1.73	5.27	0.11	3.99	0.21	28.10	0.09
VBY	21.09	0.23	26.09	1.63	5.53	0.09	8.35	0.27	25.28	0.11

Table 4. CBP benthic Monitoring Program Benthic Strata. Biotic parameters. P – Brey's secondary productivity ( $\text{gC}/\text{m}^2/\text{yr}$ ), B – biomass ( $\text{AFDW gC}/\text{m}^2$ ), A – abundance (individuals/ $\text{m}^2$ ), BIBI – benthic index of biotic integrity, S – species per sample. SE – standard error for the previous column. In parentheses – number of samples. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.

STRATUM	P	SE	B	SE	A	SE	BIBI	SE	S	SE
UPB	36.84	3.27	24.04	2.82	2,404	142	3.1	0.05	8.4	0.19
MWT	20.22	1.69	11.13	1.64	2,491	206	2.5	0.05	6.8	0.23
MET	21.17	2.50	12.27	2.80	3,150	270	2.9	0.04	9.7	0.22
PMR	19.73	2.57	18.37	3.13	1,728	176	2.1	0.06	5.6	0.25
RAP	12.90	0.74	1.72	0.44	3,607	252	2.5	0.04	9.7	0.21
PXR	8.80	0.94	3.08	1.41	1,760	129	2.5	0.05	7.5	0.22
JAM	18.09	0.97	4.81	0.77	4,396	276	2.7	0.04	11.5	0.30
MMS	20.48	3.89	2.17	0.40	8,800	2,104	2.6	0.05	10.2	0.29
YRK	16.82	1.00	2.20	0.44	4,324	288	2.6	0.04	11.7	0.25
ELR	18.65	0.69	1.61	0.23	6,035	327	2.3	0.03	13.4	0.24
VBY	14.92	0.68	2.14	0.24	4,020	270	3.1	0.04	19.3	0.42

Table 5. CBP Segments. Physico-chemical parameters. S – salinity, %SC – percent silt-clay, T – bottom water temperature (°C), DO – bottom dissolved oxygen (ppm), D – water depth (m). SE – standard error for the previous column. In parentheses – number of samples.

SEGMENT	S	SE	%SC	SE	DO	SE	D	SE	T	SE
APPTF (3)	0.1	0.0	59.8	9.8	6.6	0.4	4.4	1.1	28.8	1.2
BACOH (17)	3.1	0.6	66.8	8.5	8.4	0.7	1.6	0.2	25.1	0.5
BIGMH (22)	16.6	0.4	39.4	6.8	7.1	0.2	2.4	0.2	24.4	0.6
BOHOH (4)	5.3	1.9	65.3	17.5	8.0	0.2	2.1	0.3	24.7	1.2
BSHOH (28)	3.4	0.4	83.8	3.2	8.4	0.3	2.1	0.1	22.1	0.4
CB1TF (62)	1.1	0.2	40.2	3.8	8.1	0.2	3.3	0.3	23.5	0.3
CB2OH (113)	7.0	0.3	69.6	2.7	6.8	0.1	5.3	0.2	24.4	0.2
CB3MH (170)	11.8	0.2	66.0	2.7	4.8	0.2	7.0	0.2	24.9	0.1
CB4MH (74)	14.1	0.4	38.0	4.5	6.0	0.3	6.6	0.5	24.6	0.2
CB5MH (138)	17.0	0.2	23.4	2.7	4.8	0.2	9.8	0.4	25.4	0.2
CB6PH (62)	20.8	0.5	36.4	4.4	4.9	0.2	9.2	0.3	25.1	0.2
CB7PH (136)	22.8	0.3	21.0	2.2	5.6	0.1	9.2	0.4	24.9	0.2
CB8PH (37)	24.8	0.5	12.6	2.2	6.0	0.1	8.0	0.4	24.1	0.4
CHKOH (10)	1.5	0.6	69.1	9.1	6.7	0.4	2.0	0.3	28.7	0.5
CHOMH1 (32)	13.3	0.4	46.5	6.5	6.9	0.4	5.0	0.6	24.2	0.4
CHOMH2 (43)	10.7	0.3	48.7	5.2	6.4	0.1	4.4	0.4	23.9	0.3
CHOOH (16)	4.6	0.8	57.1	8.0	5.8	0.4	5.3	0.8	23.2	0.3
CHOTF (2)	0.8	0.1	92.9	0.9	7.2	1.1	2.9	0.1	24.7	3.0
CHSMH (81)	11.7	0.2	38.5	4.0	5.5	0.2	4.9	0.4	24.5	0.2
CHSOH (14)	6.8	0.9	74.3	6.0	6.0	0.4	3.0	0.5	24.4	0.5
CHSTF (2)	3.0	2.4	92.6	1.7	5.2	0.3	1.7	0.3	22.9	2.1
CRRMH (19)	15.6	0.4	53.8	9.8	5.1	0.3	3.2	0.5	27.4	0.5
EASMH (22)	13.4	0.4	35.8	7.8	7.0	0.3	3.8	0.6	23.8	0.3
EBEMH (44)	17.5	0.6	70.4	4.3	4.5	0.3	3.3	0.4	27.6	0.2
ELIPH (133)	21.5	0.2	51.8	3.2	5.4	0.1	6.5	0.5	27.3	0.1
ELKOH (28)	2.7	0.3	60.3	6.6	7.1	0.3	3.5	0.7	23.9	0.4
FSBMH (12)	11.8	1.1	70.0	8.3	7.2	0.4	2.3	0.2	21.0	1.0
GUNOH (33)	4.5	0.5	70.7	5.8	7.3	0.2	2.2	0.2	22.9	0.4
HNGMH (13)	15.1	0.6	37.0	7.9	7.3	0.2	2.9	0.4	25.2	0.5
JMSMH (169)	15.9	0.4	58.4	2.6	6.1	0.1	4.1	0.2	26.3	0.2
JMSOH (70)	5.8	0.5	64.6	4.7	6.3	0.2	3.8	0.3	27.8	0.2
JMSPH (44)	22.5	0.4	39.2	5.1	5.8	0.1	8.3	0.8	25.4	0.3
JMSTF (39)	0.6	0.1	71.4	5.3	6.4	0.2	4.8	0.7	26.6	0.4
LAFMH (57)	19.3	0.4	69.8	4.3	6.6	0.3	1.9	0.2	28.9	0.2
LCHMH (20)	14.9	0.4	30.3	7.8	7.9	0.3	2.6	0.5	23.7	0.4
LYNPH (2)	19.8	0.8	61.3	12.3	3.2	2.9	1.3	0.7	28.3	0.3
MAGMH (33)	11.1	0.6	53.7	6.7	4.6	0.4	3.8	0.4	24.9	0.3
MANMH (33)	15.6	0.3	41.3	6.2	7.3	0.2	2.3	0.2	25.0	0.3
MATTF (1)	2.1		16.7		6.5		2.7		25.4	
MIDOH (14)	4.4	0.6	58.9	10.5	7.6	0.2	2.1	0.2	25.3	0.5
MOBPH (38)	19.6	0.6	37.4	5.8	5.7	0.3	5.3	0.7	26.4	0.2
MPNOH (24)	6.9	1.3	48.7	8.1	4.1	0.2	6.0	0.7	27.6	0.3
MPNTF (9)	0.3	0.2	29.8	12.7	5.2	0.2	3.7	0.8	25.3	0.5

Table 5. (Continued). CBP Segments. Physico-chemical parameters. S – salinity, %SC – percent silt-clay, T – bottom water temperature (°C), DO – bottom dissolved oxygen (ppm), D – water depth (m). SE – standard error for the previous column. In parentheses – number of samples.

SEGMENT	S	SE	%SC	SE	DO	SE	D	SE	T	SE
NANMH (33)	11.0	0.6	57.9	5.9	6.8	0.2	2.8	0.2	23.5	0.4
NANOH (11)	4.2	1.1	75.8	4.7	6.4	0.7	3.7	0.8	24.4	0.6
NORTF (8)	0.6	0.2	73.8	9.0	7.0	1.3	2.6	0.2	23.6	0.5
PATMH (131)	10.2	0.3	64.5	2.7	4.3	0.2	6.0	0.3	25.9	0.1
PAXOH (28)	7.4	0.7	82.4	3.5	6.0	0.3	2.1	0.2	25.3	0.3
PAXTF (13)	0.3	0.1	77.2	6.8	6.3	0.2	2.0	0.4	22.5	0.4
PIAMH (6)	16.6	0.8	74.0	14.5	6.4	0.6	3.9	1.3	25.1	0.7
PMKOH (32)	7.1	0.9	54.9	6.5	4.8	0.2	4.6	0.4	27.8	0.3
PMKTF (10)	0.5	0.2	68.6	11.2	5.3	0.5	2.5	0.7	28.4	0.4
POCMH (19)	18.1	0.7	46.6	8.0	6.2	0.3	2.8	0.3	25.0	0.2
POCOH (11)	8.1	1.6	65.4	9.2	7.2	0.7	2.1	0.5	24.5	0.9
POCTF (1)	0.1		0.8		4.5		7.3		21.2	
POTMH (254)	13.7	0.2	61.5	2.4	4.1	0.2	7.9	0.3	25.0	0.1
POTOH (62)	4.2	0.4	80.5	2.9	6.4	0.2	4.9	0.4	23.5	0.3
POTTF (32)	0.6	0.1	79.2	4.6	7.6	0.2	5.4	0.6	22.8	0.4
RHDMH (15)	10.0	0.7	49.9	9.5	7.0	0.5	2.8	0.2	25.7	0.4
RPPMH (290)	13.4	0.2	59.8	2.3	5.2	0.1	5.4	0.3	26.1	0.1
RPPOH (16)	2.8	0.5	82.6	5.7	6.2	0.3	2.6	0.6	26.2	0.5
RPPTF (29)	0.4	0.1	71.9	5.3	7.0	0.3	2.4	0.4	26.7	0.4
SASOH (23)	2.9	0.4	66.2	7.1	8.0	0.3	3.2	0.3	23.1	0.5
SBEMH (134)	19.9	0.3	53.7	2.7	4.4	0.2	3.9	0.4	28.5	0.1
SEVMH (35)	11.0	0.7	57.8	5.4	3.8	0.5	4.8	0.4	25.0	0.4
SOUHM (28)	10.6	0.7	42.0	6.9	5.1	0.4	3.2	0.3	25.2	0.5
TANMH (103)	16.6	0.2	24.2	3.2	6.7	0.1	5.5	0.4	25.4	0.2
WBEMH (52)	21.0	0.5	70.6	4.9	6.2	0.3	1.7	0.2	28.6	0.3
WICMH (26)	11.6	0.6	52.4	6.8	7.0	0.2	3.1	0.3	24.0	0.4
WSTMH (15)	12.1	0.5	74.0	8.3	6.3	0.4	2.5	0.3	24.3	0.5
YRKMH (179)	17.0	0.3	64.9	2.4	5.5	0.1	3.5	0.2	27.5	0.1
YRKPH (91)	20.7	0.3	55.0	3.8	4.6	0.2	7.2	0.6	26.7	0.1

Table 6. CBP Segments. Biotic parameters. P – Brey's secondary productivity (gC/m<sup>2</sup>/yr), B – biomass (AFDW/gC/m<sup>2</sup>), A – abundance (individual/m<sup>2</sup>), BIBI – benthic index of biotic integrity, S – species per sample. SE – standard error for the previous column. In parentheses – number of samples.

SEGMENTS	P	SE	B	SE	A	SE	B_IBI	SE	S	SE
APPTF (3)	17.83	3.18	0.7	0.10	4,636	841	2.3	0.33	7.67	0.33
BACOH (17)	8.86	3.89	1.1	0.37	4,511	1,929	2.2	0.16	5.53	0.67
BIGMH (22)	6.53	1.20	0.8	0.18	2,784	524	3.2	0.16	12.23	1.30
BOHOH (4)	69.69	39.42	58.6	41.02	2,079	422	3.1	0.49	6.00	1.47
BSHOH (28)	27.73	7.60	24.4	8.65	2,292	682	2.6	0.13	5.82	0.54
CB1TF (62)	48.64	10.44	38.3	9.71	3,489	371	3.2	0.11	9.37	0.44
CB2OH (113)	64.62	7.14	44.5	5.92	2,447	218	3.5	0.06	9.35	0.26
CB3MH (170)	15.65	1.59	6.0	1.30	1,960	202	2.9	0.08	7.55	0.29
CB4MH (74)	58.19	16.61	5.7	1.63	24,505	8,336	2.3	0.11	8.28	0.52
CB5MH (138)	6.23	0.59	1.1	0.37	1,658	130	2.6	0.08	11.32	0.60
CB6PH (62)	15.50	1.50	2.8	0.70	3,746	305	3.3	0.10	19.81	1.00
CB7PH (136)	15.25	0.95	1.9	0.16	4,761	633	3.3	0.06	21.48	0.65
CB8PH (37)	19.97	3.35	3.9	1.29	5,473	810	3.3	0.12	24.03	1.42
CHKOH (10)	21.94	6.19	8.8	4.06	2,491	326	3.4	0.21	8.70	0.63
CHOMH1 (32)	8.73	1.41	1.0	0.13	3,894	1,726	2.4	0.14	9.09	0.76
CHOMH2 (43)	17.43	2.16	2.7	0.37	6,834	2,084	3.0	0.11	11.40	0.35
CHOOH (16)	18.54	5.05	12.2	5.10	2,228	533	3.0	0.26	8.13	0.95
CHOTF (2)	19.90	19.59	26.8	26.77	897	829	2.8	0.20	4.50	2.50
CHSMH (81)	14.68	1.16	2.8	0.48	3,194	450	2.8	0.10	9.77	0.44
CHSOH (14)	101.38	34.73	109.0	40.25	2,197	798	3.0	0.16	9.14	0.97
CHSTF (2)	34.49	28.80	41.0	40.82	4,295	1	1.7	0.30	9.00	3.00
CRRMH (19)	9.40	2.28	0.6	0.11	2,066	544	2.0	0.17	9.05	1.06
EASMH (22)	39.29	18.31	4.7	2.46	25,737	12,679	2.1	0.16	8.64	1.04
EBEMH (44)	21.27	2.11	2.2	0.73	6,999	962	2.2	0.09	13.43	0.62
ELIPH (133)	16.56	0.89	2.3	0.63	4,088	287	2.6	0.07	15.46	0.52
ELKOH (28)	42.10	11.24	26.8	8.00	2,153	434	2.9	0.15	7.68	0.75
FSBMH (12)	6.81	1.39	1.4	0.40	1,651	281	3.3	0.28	10.50	0.92
GUNOH (33)	45.69	7.02	54.2	10.90	2,002	509	3.0	0.10	7.88	0.50
HNGMH (13)	7.19	2.06	0.7	0.18	2,297	509	2.8	0.14	11.69	0.87
JMSMH (169)	13.94	0.80	1.6	0.22	4,275	438	2.7	0.05	11.59	0.39
JMSOH (70)	22.45	3.21	6.3	2.04	4,711	654	2.6	0.09	8.30	0.24
JMSPH (44)	16.66	1.94	7.3	3.07	3,535	326	3.1	0.12	19.07	1.10
JMSTF (39)	28.23	4.13	14.5	4.26	4,569	617	2.8	0.15	8.59	0.40
LAFMH (57)	17.88	1.05	1.1	0.10	5,854	591	2.4	0.08	13.28	0.53
LCHMH (20)	26.98	9.10	3.0	1.02	8,956	3,151	2.5	0.14	10.05	0.52
LYNPH (2)	39.80	13.80	3.8	1.77	4,955	1,273	2.3	0.33	11.00	3.00
MAGMH (33)	10.29	2.67	1.4	0.32	2,058	427	2.3	0.16	6.85	0.81
MANMH (33)	5.68	0.75	0.7	0.15	1,794	229	2.9	0.15	10.52	0.80
MATTF (1)	118.33		122.7		4,499		1.7		11.00	
MIDOH (14)	73.69	22.23	49.4	14.50	1,592	477	3.2	0.14	6.93	0.83
MOBPH (38)	15.45	1.49	1.3	0.17	3,772	314	2.7	0.12	17.24	1.12
MPNOH (24)	11.63	1.70	1.0	0.17	3,382	794	2.9	0.19	7.29	0.67
MPNTF (9)	32.80	21.68	14.0	11.20	2,558	561	3.2	0.22	7.33	0.69

Table 6 (Continued). CBP Segments. Biotic parameters. P – Brey’s secondary productivity (gC/m<sup>2</sup>/yr), B – biomass (AFDW/gC/m<sup>2</sup>), A – abundance (individual/m<sup>2</sup>), BIBI – benthic index of biotic integrity, S – species per sample. SE – standard error for the previous column. In parentheses – number of samples.

SEGMENTS	P	SE	B	SE	A	SE	B_IBI	SE	S	SE
NANMH (33)	7.71	1.29	1.7	0.73	2,498	495	3.0	0.13	10.30	0.46
NANOH (11)	48.79	35.57	64.4	57.94	2,878	776	3.1	0.28	7.45	0.77
NORTF (8)	2.76	1.56	0.4	0.17	4,085	1,591	2.6	0.40	7.25	0.96
PATMH (131)	15.40	1.74	3.9	0.55	2,265	343	2.3	0.10	6.01	0.40
PAXMH (309)	7.69	0.60	1.1	0.10	1,649	124	2.5	0.05	7.51	0.24
PAXOH (28)	24.19	9.39	26.0	17.34	2,573	812	2.9	0.20	7.14	0.59
PAXTF (13)	1.46	0.52	0.7	0.24	2,639	575	2.3	0.23	7.85	1.12
PIAMH (6)	4.18	0.61	0.4	0.04	1,008	205	1.9	0.14	5.83	0.65
PMKOH (32)	15.42	3.30	2.4	1.27	5,707	1,914	2.9	0.18	7.53	0.50
PMKTF (10)	33.57	13.63	17.6	9.63	2,218	670	3.3	0.21	7.10	0.86
POCMH (19)	7.59	1.36	0.7	0.14	2,177	454	2.7	0.22	13.32	1.75
POCOH (11)	24.40	7.59	2.8	0.93	6,444	1,530	2.7	0.16	11.55	1.60
POCTF (1)	0.82		0.1		659		2.5		5.00	
POTMH (254)	4.52	0.78	2.1	1.04	1,154	189	1.8	0.05	4.47	0.30
POTOH (62)	55.24	7.16	52.4	7.05	2,149	269	3.1	0.10	8.05	0.35
POTTF (32)	69.15	18.75	78.5	26.05	5,408	849	2.9	0.19	9.09	0.63
RHDMH (15)	19.39	4.23	1.9	0.43	3,820	794	2.7	0.25	9.33	0.56
RPPMH (290)	11.45	0.61	1.0	0.15	3,511	278	2.5	0.05	10.01	0.24
RPPOH (16)	17.31	5.12	9.7	7.00	3,517	781	2.7	0.29	7.88	0.91
RPPTF (29)	27.19	5.04	4.9	3.36	5,568	1,120	3.3	0.18	9.00	0.47
SASOH (23)	24.58	13.63	10.8	3.96	2,617	1,064	2.7	0.13	6.91	0.66
SBEMH (134)	21.07	1.59	1.1	0.08	8,280	821	2.0	0.05	12.00	0.34
SEVMH (35)	5.94	1.13	0.9	0.22	1,502	359	2.2	0.17	6.00	0.81
SOUMH (28)	14.47	2.59	1.3	0.23	4,024	726	2.3	0.18	9.07	0.81
TANMH (103)	7.28	0.53	0.8	0.06	1,965	120	3.1	0.07	13.90	0.51
WBEMH (52)	16.56	1.67	0.9	0.06	4,903	481	2.3	0.07	11.87	0.37
WICMH (26)	15.10	5.45	1.3	0.31	4,916	1,384	3.0	0.18	10.92	0.73
WSTMH (15)	21.85	7.50	3.2	1.30	3,453	1,346	3.0	0.21	8.67	0.93
YRKMH (179)	17.63	1.18	1.2	0.07	5,127	408	2.5	0.05	12.02	0.26
YRKPH (91)	13.79	0.96	1.5	0.33	2,937	204	2.5	0.08	14.44	0.52

Table 7. Top twenty secondary production dominant species for CBP segments JMSMH and SBEMH for the period 1996 through 2009. Values shown are segment mean values for each species. A – amphipod, B – bivalve, C – cumacean, G – gastropod, H – hemichordate, I – isopod, In – insect, N – nemertine, O – oligochaete, P – polychaete, Ph – phoronid.

Segment JMSMH		Segment SBEMH	
	Mean Brey's Productivition (g C/m <sup>2</sup> /yr)		Mean Brey's Productivition (g C/m <sup>2</sup> /yr)
<i>Mediomastus ambiseta</i> (P)	2.20	<i>Mediomastus ambiseta</i> (P)	6.01
<i>Neanthes succinea</i> (P)	1.45	<i>Streblospio benedicti</i> (P)	3.31
<i>Leptocheirus plumulosus</i> (A)	1.44	<i>Leitoscoloplos</i> spp. (P)	2.06
<i>Paraprionospio pinnata</i> (P)	0.94	<i>Laeonereis culveri</i> (P)	1.36
<i>Macoma balthica</i> (B)	0.93	<i>Heteromastus filiformis</i> (P)	1.06
<i>Streblospio benedicti</i> (P)	0.64	<i>Paraprionospio pinnata</i> (P)	1.00
<i>Heteromastus filiformis</i> (P)	0.59	<i>Cyathura polita</i> (I)	0.80
<i>Glycinde solitaria</i> (P)	0.55	<i>Glycinde solitaria</i> (P)	0.67
<i>Tubificoides heterochaetus</i> (O)	0.52	<i>Tubificoides</i> spp. (O)	0.43
<i>Leitoscoloplos</i> spp. (P)	0.42	<i>Neanthes succinea</i> (P)	0.40
<i>Cyathura polita</i> (I)	0.35	<i>Eteone heteropoda</i> (P)	0.36
<i>Tubificoides</i> spp. (O)	0.29	<i>Tubificoides heterochaetus</i> (O)	0.35
<i>Marenzelleria viridis</i> (P)	0.26	<i>Loimia medusa</i> (P)	0.27
<i>Macoma mitchelli</i> (B)	0.26	<i>Capitella capitata</i> complex(P)	0.22
<i>Leucon americanus</i> (C)	0.20	<i>Leucon americanus</i> (C)	0.22
<i>Mercenaria mercenaria</i> (B)	0.19	Nemertina spp. (N)	0.19
<i>Loimia medusa</i> (P)	0.18	<i>Tharyx</i> sp. A Morris (P)	0.17
Nemertina spp. (N)	0.17	<i>Macoma mitchelli</i> (B)	0.17
<i>Polypedilum</i> spp. (In)	0.15	Oligochaeta spp. (O)	0.16
<i>Rangia cuneata</i> (B)	0.14	<i>Cyclaspis varians</i> (C)	0.15

Table 8. Top twenty density dominant species for CBP segments JMSMH and SBEMH for the period 1996 through 2009. Values shown are segment mean values for each species. A – amphipod, B – bivalve, C – cumacean, G – gastropod, H – hemichordate, I – isopod, In – insect, N – nemertine, O – oligochaete, P – polychaete, Ph – phoronid.

Segment JMSMH		Segment SBEMH	
	Mean Density (#/m <sup>2</sup> )		Mean Density (#/m <sup>2</sup> )
<i>Mediomastus ambiseta</i> (P)	1330	<i>Mediomastus ambiseta</i> (P)	3859
<i>Tubificoides heterochaetus</i> (O)	589	<i>Streblospio benedicti</i> (P)	2259
<i>Streblospio benedicti</i> (P)	526	<i>Tubificoides heterochaetus</i> (O)	313
<i>Leptocheirus plumulosus</i> (A)	519	<i>Tubificoides</i> spp. (O)	281
<i>Tubificoides</i> spp. (O)	203	<i>Laeonereis culveri</i> (P)	206
<i>Glycinde solitaria</i> (P)	135	<i>Paraprionospio pinnata</i> (P)	161
<i>Paraprionospio pinnata</i> (P)	134	<i>Oligochaeta</i> spp. (O)	155
<i>Neanthes succinea</i> (P)	128	<i>Leitoscoloplos</i> spp (P)	149
<i>Leucon americanus</i> (C)	85	<i>Glycinde solitaria</i> (P)	113
<i>Heteromastus filiformis</i> (P)	82	<i>Heteromastus filiformis</i> (P)	104
<i>Acteocina canaliculata</i> (G)	42	<i>Cyathura polita</i> (I)	90
<i>Macoma balthica</i> (B)	39	<i>Leucon americanus</i>	75
Nemertina (N)	36	<i>Capitella capitata complex</i> (P)	64
<i>Cyathura polita</i> (I)	34	<i>Eteone heteropoda</i> (P)	58
<i>Leitoscoloplos</i> spp. (P)	32	Nemertina (N)	36
<i>Ampelisca</i> spp. (A)	28	<i>Tharyx</i> sp. A Morris (P)	34
<i>Polydora cornuta</i> (P)	28	<i>Neanthes succinea</i> (P)	33
<i>Marenzelleria viridis</i> (P)	28	<i>Cyclaspis varians</i> (C)	30
<i>Phoronis</i> spp. (Ph)	23	<i>Hobsonia florida</i> (P)	29
<i>Macoma mitchelli</i> (B)	22	Tubificidae spp. (O)	28

Table 9. Top twenty biomass dominant species for CBP segments JMSMH and SBEMH for the period 1996 through 2009. Values shown are segment mean values for each species. A – amphipod, B – bivalve, C – cumacean, G – gastropod, H – hemichordate, I – isopod, In – insect, N – nemertine, O – oligochaete, P – polychaete, Ph – phoronid.

Segment JMSMH		Segment SBEMH	
Taxon	Mean Biomass (g C/m <sup>2</sup> )	Taxon	Mean Biomass (g C/m <sup>2</sup> )
<i>Mercenaria mercenaria</i> (B)	0.259	<i>Leitoscoloplos</i> spp. (P)	0.198
<i>Macoma balthica</i> (B)	0.223	<i>Mediomastus ambiseta</i> (P)	0.131
<i>Neanthes succinea</i> (P)	0.129	<i>Heteromastus filiformis</i> (P)	0.085
<i>Paraprionospio pinnata</i> (P)	0.075	<i>Laeonereis culveri</i> (P)	0.071
<i>Mediomastus ambiseta</i> (P)	0.066	<i>Streblospio benedicti</i> (P)	0.069
<i>Leptocheirus plumulosus</i> (A)	0.056	<i>Paraprionospio pinnata</i> (P)	0.065
<i>Macoma mitchelli</i> (B)	0.051	<i>Cyathura polita</i> (I)	0.052
<i>Polinices duplicatus</i> (G)	0.050	<i>Loimia medusa</i> (P)	0.041
<i>Rangia cuneata</i> (B)	0.047	<i>Neanthes succinea</i> (P)	0.035
<i>Leitoscoloplos</i> spp. (P)	0.046	<i>Glycinde solitaria</i> (P)	0.034
<i>Heteromastus filiformis</i> (P)	0.043	<i>Glycera Americana</i> (P)	0.033
<i>Glycinde solitaria</i> (P)	0.032	<i>Macoma mitchelli</i> (B)	0.033
<i>Cyathura polita</i> (I)	0.029	<i>Macoma balthica</i> (B)	0.024
<i>Mya arenaria</i> (B)	0.029	<i>Nemertina</i> (N)	0.024
<i>Glycera Americana</i> (P)	0.028	<i>Eteone heteropoda</i> (P)	0.018
<i>Loimia medusa</i> (P)	0.026	<i>Tagelus plebeius</i> (B)	0.016
<i>Marenzelleria viridis</i> (P)	0.024	<i>Hemichordata</i> (H)	0.014
<i>Streblospio benedicti</i> (P)	0.024	<i>Tubificoides</i> spp. (O)	0.013






Figure 1. Benthic Secondary Productivity as a function of Total Community Biomass comparing Edgar's equation versus Brey's. Equation applied for each species and summed.




Figure 2. Benthic Secondary Productivity as a function of Total Community Biomass using the Edgar equation. Equation applied for each species and summed.




Figure 3. Benthic Secondary Productivity as a function of Total Community Biomass using the Brey equation. Equation applied for each species and summed.

● BREY01\_PRODUCTION\_GC    ◆ EDGAR\_SPECIES\_LEVEL1





Figure 4. Benthic Secondary Productivity as a function of Total Community Biomass comparing Edgar's equation versus Brey's. Equation applied for each species and summed. A. Biomass values to 100 gC/m<sup>2</sup> Biomass. B. Biomass axis limited to  $\leq 5$  gC/m<sup>2</sup>/yr.




Figure 5. Mean secondary production ( $\text{gC/m}^2/\text{yr}$ ) by the habitat types of Weisberg et al. (1997). Bar indicates one standard error. All random data from 1996 -2009  $n = 3,919$ .




Figure 6. Mean standing stock biomass ( $\text{gC/m}^2$ ) by the habitat types of Weisberg et al. (1997). Bar indicates one standard error. All random data from 1996 -2009  $n = 3,919$ .




Figure 7. Mean community abundance by the habitat types of Weisberg et al. 1997). Bar indicates one standard error. All random data from 1996 -2009 n = 3,919.




Figure 8. Mean B-IBI by the habitat types of Weisberg et al. 1997). Bar indicates one standard error. All random data from 1996 -2009 n = 3,919.




Figure 9. Mean species per sample by the habitat types of Weisberg et al. 1997). Bar indicates one standard error. All random data from 1996 -2009 n = 3,919.




Figure 10. Mean Shannon Index by the habitat types of Weisberg et al. 1997). Bar indicates one standard error. All random data from 1996 -2009 n = 3,919.




Figure 11. Chesapeake Bay Benthic Monitoring Strata






Figure 12. Mean secondary production (gC/m<sup>2</sup>/yr) by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.




Figure 13. Mean standing stock biomass (gC/m<sup>2</sup>) by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.




Figure 14. Mean community abundance by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.




Figure 15. Mean B-IBI by the habitat types by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.




Figure 16. Mean species per sample ) by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.




Figure 17. Mean Shannon Index ) by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.




Figure 18. Log<sub>10</sub> mean secondary production (gC/m<sup>2</sup>/yr) by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.




Figure 19. Log<sub>10</sub> mean standing stock biomass (gC/m<sup>2</sup>) by the CBP benthic strata. Bar indicates one standard error. UPB – Upper Mainstem, MWT – Maryland Western Tributaries, MET – Maryland Eastern Tributaries, PMR – Potomac River, RAP – Rappahannock River, PXR – Patuxent River, JAM – James River, MMS – Middle Mainstem, YRK – York River, ELR – Elizabeth River, VBY – Virginia Mainstem.



Figure 20. Chesapeake Bay Program segments.




Figure 21. Chesapeake Bay Program segments showing levels of benthic secondary productivity (see insert at top).




Figure 22. CBP segments. Shown are secondary productivity values in  $\text{gC}/\text{m}^2/\text{yr}$ . Colors correspond to the intervals of Figure 21.




Figure 23. Composite Chesapeake Bay summer DO concentration from 1996 to 2004. Large and small circles represent sample sites. Shading and dot color denotes DO concentration as stated in Figure key.






Figure 24. CB4MH secondary productivity values in  $\text{gC/m}^2/\text{yr}$ . A. all water depths. B. Depth  $\geq 40$  m.




Figure 25. CB4MH secondary productivity values in  $\text{gC/m}^2/\text{yr}$ . A.. Depth  $\geq 20$  m.  
B. Depth  $\geq 10$  m.


**Lower Potomac River (POTMH)**

Figure 26. POTMH secondary productivity values in gC/m<sup>2</sup>/yr. Productivity is shown on a log scale due to the wide range of values. Insert shows average secondary productivity by water depth intervals and the percentage of samples with no benthos.




Figure 27. CBP segments and low dissolved oxygen. Shown are secondary productivity values in gC/m<sup>2</sup>/yr. Colors correspond to the intervals of Figure 21. Segments with no color and cross hatching have mean summer dissolved oxygen levels <5 mg/l.




Figure 28. CBP segments and high urban land-use. Shown are secondary productivity values in  $\text{gC}/\text{m}^2/\text{yr}$ . Colors correspond to the intervals of Figure 21. Segments with no color and cross hatching are located in either the Maryland Western Tributaries benthic stratum or the Elizabeth River watershed.




Figure 29. CBP segments and high urban land-use (the Maryland Western Tributaries benthic stratum or the Elizabeth River watershed). Shown are secondary productivity values in gC/m<sup>2</sup>/y compared to biomass in gC/m<sup>2</sup>. Colored vertical lines s correspond to the intervals of Figure 21. Green circle indicates SBEMH from the Elizabeth River.




Figure 30. Mean secondary production (gC/m<sup>2</sup>/yr) comparing JMSMH (James River mesohaline), POTMH (lower Potomac River mesohaline), PATMH (Patapsco River) and SMEMH (Southern Branch of the Elizabeth River).




Figure 31. Mean standing stock biomass (gC/m<sup>2</sup>) comparing JMSMH (James River mesohaline), POTMH (lower Potomac River mesohaline), PATMH (Patapsco River) and SMEMH (Southern Branch of the Elizabeth River).




Figure 32. Mean community abundance comparing JMSMH (James River mesohaline), POTMH (lower Potomac River mesohaline), PATMH (Patapsco River) and SBEMH (Southern Branch of the Elizabeth River).




Figure 33. Mean B-IBI comparing JMSMH (James River mesohaline), POTMH (lower Potomac River mesohaline), PATMH (Patapsco River) and SBEMH (Southern Branch of the Elizabeth River).






Figure 34. Mean species per sample comparing JMSMH (James River mesohaline), POTMH (lower Potomac River mesohaline), PATMH (Patapsco River) and SBEMH (Southern Branch of the Elizabeth River).




Figure 35. Mean Shannon Index comparing JMSMH (James River mesohaline), POTMH (lower Potomac River mesohaline), PATMH (Patapsco River) and SBEMH (Southern Branch of the Elizabeth River).




Figure 36. CBP segment secondary productivity values in gC/m<sup>2</sup>/yr and the B-IBI. A. All segments. B. Segments in the two highest productivity categories > 20 and >40 gC/m<sup>2</sup>/yr

Appendix A. Conversion factors used to estimate energetic content of ash free dry weight (AFDW) of individual taxa. An \* indicates a higher level taxonomic group conversion factor was used for the group (e.g. Mollusca for Scaphopoda). A \*\* indicates the general Annelida conversion factor was applied. A † indicates the conversion factor for Crustacea was applied. A ‡ indicates the conversion factor for Ascidiacea was applied.

<b>Phylogenetic Group</b>	<b>Conversion</b>
Anthozoa	24.46
Annelida	23.33
Mollusca	23.04
Gastropoda	23.63
Polyplacophora	23.27
Bivalvia	22.79
Scaphopoda*	23.04
Cephalopoda	22.03
Arthropoda	
Crustacea	22.57
Amphipoda	22.74
Cephalocarida*	22.57
Cumacea	22.74
Decapoda	22.26
Isopoda*	22.74
Mysidacea	23.00
Stomatopoda	22.57
Tanaidacea*	22.57
Merostomata†	22.57
Insecta (General)	23.81
Coleoptera	23.81
Collembola	23.81
Chironomidae	23.44
Diptera	23.81
Ephemeroptera	26.07
Megaloptera	23.81
Odonata	23.65
Plecoptera	23.81
Trichoptera	24.12
Echinodermata	
Asteroidea	20.81
Ophiuroidea	21.75
Echinoidea	20.53
Holothuroidea	22.95
Chordata	
Ascidiacea	19.01
Cephalochordata‡	19.01
Miscellaneous groups	
Nemertina**	23.33
Echiurida**	23.33
Phoronida**	23.33
Sipuncula**	23.33

Appendix B. Comparison of benthic secondary productivity BCS of this study to Diaz and Schaffner (1990) (D & S). For euhaline habitat the D&S value was used for both the Brey and Edgar estimates of our study.

Habitat Type and Area			Diaz & Schaffner (1990)		Present Study Area weighed BSC		Total Habitat Productivity (metric tons of C/yr)		
Major habitat	Area km <sup>2</sup>	% of Habitat	Mean BSC	Area weighted BSC	ODU Edgar	ODU Brey	D & S	ODU Brey's	ODU Edgar's
Tidal Freshwater									
Mud	455	0.80	1.8	1.44					
Sand	102	0.18	145.5	26.17					
Mixed	10	0.02	289.2	5.10					
<b><u>Total area</u></b>	<b><u>567</u></b>			<b><u>32.72</u></b>	<b><u>126.28</u></b>	<b><u>32.44</u></b>	<b><u>18,552</u></b>	<b><u>18,396</u></b>	<b><u>71,603</u></b>
Oligohaline									
Mud	496	0.79	14.4	11.32					
Sand	59	0.09	18.0	1.68					
Mixed	76	0.12	21.7	2.61					
<b><u>Total area</u></b>	<b><u>631</u></b>			<b><u>15.62</u></b>	<b><u>151.11</u></b>	<b><u>36.00</u></b>	<b><u>9,854</u></b>	<b><u>22,714</u></b>	<b><u>95,351</u></b>
Low Mesohaline									
Mud	393	0.75	14.4	10.74					
Sand	98	0.19	41.0	7.62					
Mixed	36	0.06	10.8	0.74					
<b><u>Total area</u></b>	<b><u>527</u></b>			<b><u>19.10</u></b>	<b><u>81.81</u></b>	<b><u>24.55</u></b>	<b><u>10,066</u></b>	<b><u>12,937</u></b>	<b><u>43,113</u></b>
High Mesohaline									
Mud	1525	0.48	8.1	3.88					
Sand	1388	0.44	8.8	3.84					
Mixed	268	0.08	25.0	2.11					
<b><u>Total area</u></b>	<b><u>3181</u></b>			<b><u>9.83</u></b>	<b><u>16.00</u></b>	<b><u>12.00</u></b>	<b><u>31,267</u></b>	<b><u>38,172</u></b>	<b><u>50,896</u></b>
Polyhaline									
Mud	509	0.18	9.0	1.64					
Sand	1764	0.63	32.0	20.27					
Mixed	512	0.18	15.6	2.87					
<b><u>Total area</u></b>	<b><u>2785</u></b>			<b><u>24.78</u></b>	<b><u>22.00</u></b>	<b><u>15</u></b>	<b><u>69,016</u></b>	<b><u>41,775</u></b>	<b><u>61,270</u></b>
Euhaline									
Mud	148	0.16	17.2	2.69					
Sand	768	0.81	5.7	4.63					
Mixed	29	0.03	28.6	0.88					
<b><u>Total area</u></b>	<b><u>945</u></b>			<b><u>8.20</u></b>			<b><u>7,753</u></b>	<b><u>7,753</u></b>	<b><u>7,753</u></b>
<b><u>Total for Chesapeake Bay</u></b>							<b><u>146,507</u></b>	<b><u>141,746</u></b>	<b><u>329,986</u></b>