

Climate Change Impacts for Chesapeake Bay Fish Production

Modeled with the


Chesapeake Atlantis Model

Tom Ihde – ERT, Inc.

Tom.Ihde@noaa.gov

Howard Townsend – NMFS

SF-GIT – 1 June, 2015




ASSESSMENT

MANAGEMENT



Climate Change Assumptions





- 50 years from now:
 - ✓ Increased water temperature (1.5°C)
 - ✓ Salinity (+/- 2 ppt)




Habitat Scenario Assumptions

- 50% loss of Marsh (area & biomass)
 Due to multiple, interacting factors:
 - shoreline armoring
 - subsidence
 - sea level rise
- 50% loss of Seagrass



Sensitivity To Climate Change Selected Group Effects of Interest to Management



BA – Bay Anchovy

SB - Striped Bass

AM – Atlantic Menhaden

SF – Summer Flounder


BC - Blue Crab

FF – All Finfish

KF – Key Forage



Sensitivity To Climate Change Selected Group Effects of Interest to Management






Current Conditions

Striped Bass

Temperature increase & Habitat Loss



Next...

- ✓ Animal preferences for temperature and salinity
 establish new 'base' model
- ✓ Rerun all models to date, ensures apple-apple comparisons with new base model
- ✓ Interesting combinations what is most useful?
- ✓ Add TMDL scenario to comparison/ combinations

Thanks to:











Marine and Atmospheric Research





Northeast Fisheries Science Center







Current Conditions

Blue Crab

Temperature increase & Habitat Loss



SAV Modeled in CAM



Marsh Modeled in CAM

