RELEVANT REFERENCES

Shell Assimilation: Quantile Regression

Higgins, C. B., Stephenson, K., and Brown, B. L. 2011. Nutrient Bioassimilation Capacity of Aquacultured Oysters: Quantification of an Ecosystem Service. *Journal of Environment Quality*, 40 (1), 271.

Higgins unpubl. data (Choptank River and Lynnhaven River)

Kellogg et al. unpubl. data (Harris Creek)

Parker and Bricker unpubl. data (Chesapeake Bay main stem, Chester River, Honga River, and Potomac River)

Powell, E. N., Mann, R., Ashton-Alcox, K. A., Kim, Y., and Bushek, D. 2016. The allometry of oysters: spatial and temporal variation in the length–biomass relationships for Crassostrea virginica. *Journal of the Marine Biological Association of the United Kingdom*, *96*(5), 1127–1144.

Luckenbach, M. W., and Ross, P. G. 2009. Recruitment, Substrate Quality and Standing Stock Monitoring in Support of NOAA-ACOA Oyster Restoration Projects in the Great Wicomico, Rappahannock, Piankatank and Lynnhaven River Basins, 2004-2006: Supplementary Materials. Eastern Shore Laboratory, Virginia Institute of Marine Science.

Shell Assimilation: % Nitrogen and Phosphorus Content

Grizzle, R. E., Ward, K. M., Peter, C. R., Cantwell, M., Katz, D., and Sullivan, J. 2016. Growth, morphometrics and nutrient content of farmed eastern oysters, Crassostrea virginica (Gmelin), in New Hampshire, USA. *Aquaculture Research*, 1–13.

Higgins, C. B., Stephenson, K., and Brown, B. L. 2011. Nutrient Bioassimilation Capacity of Aquacultured Oysters: Quantification of an Ecosystem Service. *Journal of Environment Quality*, 40 (1), 271.

Kellogg, M. L., Cornwell J. C., Owens, M. S. and K. T. Paynter. 2013. Denitrification and nutrient assimilation on a restored oyster reef. *Marine Ecology Progress Series* 480:1-19.

Kellogg unpubl. data (Choptank River)

Reitsma, J., Murphy, D.C., Archer, A.F., and York, R.H. 2016. Nitrogen extraction potential of wild and cultured bivalves harvested from nearshore waters of Cape Cod, USA. Marine Pollution Bulletin (2016), http://dx.doi.org/10.1016/j.marpolbul.2016.12.072

Sebastiano, D., Levinton, J. S., Doall, M., and Kamath, S. 2015. Using a shellfish harvest strategy to extract high nitrogen inputs in urban and suburban coastal bays: practical and economic implications. *Journal of Shellfish Research*, 34(2), 573-583.

Shell Assimilation: Shell Dissolution-Related

Christmas, J.F., M.R. McGinty, D.A. Randle, and S.J. Jordan. 1997. Oyster shell disarticulation in three Chesapeake Bay tributaries. Journal of Shellfish Research 16(1): 115-123.DePiper et al. 2016

DePiper, G.S., D.W. Lipton, and R.N. Lipcius. 2016. Valuing Ecosystem Services: Oysters, Denitrification, and Nutrient Trading Programs. Marine Resource Economics 32(1).

Jordan-Cooley, W. C., R. N. Lipcius, L. B. Shaw, J. Shen, and J. Shi. 2011. "Bistability in a Differential Equation Model of Oyster Reef Height and Sediment Accumulation." Journal of Theoretical Biology 289:1–11.

Mann, R., M. Southworth, J.M. Harding, and J.A. Wesson. 2009. Population Studies of the Native Easter Oyster Crassostra virginica (Gmelin, 1791) in the James River, Virginia, USA. Journal of Shellfish Research 28(2): 193-220.

Powell, E.N. and J.M. Klinck. 2007. Is oyster shell a sustainable estuarine resource? Journal of Shellfish Research 26(1): 181-194

Powell, E.N., J.N. Kraeuter, K.A. Ashton-Alcox. 2006. How long does oyster shell last on an oyster reef? Estuarine, Coastal and Shelf Science 69: 531-542.Smith G.F., D. G. Bruce, E. B. Roach, A. Hansen, R. I. E. Newell & A. M. McManus (2005) Assessment of Recent Habitat Conditions of Eastern Oyster Crassostrea virginica Bars in Mesohaline Chesapeake Bay. North American Journal of Fisheries Management 25(4), 1569-1590. DOI: 10.1577/M04-058.1Waldbusser et al. 2011

Smith G.F., D. G. Bruce, E. B. Roach, A. Hansen, R. I. E. Newell & A. M. McManus (2005) Assessment of Recent Habitat Conditions of Eastern Oyster Crassostrea virginica Bars in Mesohaline Chesapeake Bay. North American Journal of Fisheries Management 25(4), 1569-1590. DOI: 10.1577/M04-058.1

Waldbusser, G.G., R.A. Steenson, and M.A. Green. 2011. Oyster shell dissolution rates in estuarine waters: effects of pH and shell legacy. Journal of Shellfish Research 30(3): 659-669.

Wilberg, M.J., M.E. Livings, J.S. Barkman, B.T. Morris, and J.M. Robinson. 2011. Overfishing, disease, habitat loss, and potential extirpation of oysters in upper Chesapeake Bay. Marine Ecology Progress Series 436: 131-144.

Enhanced Denitrification

Higgins, C.B., C. Tobias, M.F. Piehler, A.R. Smyth, R.F. Dame, K. Stephenson, and B.L. Brown. 2013. Effect of aquacultured oyster biodeposition on sediment N2 production in Chesapeake Bay. Marine Ecology Progress Series 473: 7–27.

Hoellein, T. J., Zarnoch, C. B., and Grizzle, R. E. 2014. Eastern oyster (Crassostrea virginica) filtration, biodeposition, and sediment nitrogen cycling at two oyster reefs with contrasting water quality in Great Bay Estuary (New Hampshire, USA. Biogeochemistry 122: 113-129.

Holyoke, R. R. 2008. Biodeposition and biogeochemical processes in shallow, mesohaline sediment of Chesapeake Bay. Ph.D., University of Maryland, College Park, MD.

Humphries et al. 2016. Directly Measured Denitrification Reveals Oyster Aquaculture and restored Oyster Reefs Remove Nitrogen at Comparable High Rates. Frontiers in Marine Science 3(74): 1-10, http://journal.frontiersin.org/article/10.3389/fmars.2016.00074/full.

Kellogg, M. L., Cornwell J. C., Owens, M. S. and K. T. Paynter. 2013. Denitrification and nutrient assimilation on a restored oyster reef. *Marine Ecology Progress Series* 480:1-19.

Lunstrum. A. 2015. Oyster aquaculture impacts on sediment nitrogen cycling and efficacy as a nutrient bioextraction tool in a tributary of Chesapeake Bay. Master's Thesis. University of Virginia.

Piehler M.F. and A.R. Smyth. 2011. Habitat-specific distinctions in estuarine denitrification affect both ecosystem function and services. Ecosphere 2: art12. doi:10.1890/ES10-00082.1.

Sisson M., M. L.Kellogg, M.W. Luckenbach, R. Lipcius, A. Colden, J.C. Cornwell, and M.S. Owens. 2011. Assessment of oyster reefs in Lynnhaven River as a Chesapeake Bay TMDL Best Management Practice. Final Report to the U. S. Army Corps of Engineers, Norfolk District and the City of Virginia Beach, Special Report No. 429 in Applied Marine Science and Ocean Engineering, Virginia Institute of Marine Science, Gloucester Point, Virginia. Pp.

Smyth, A. R., N. R. Geraldi, and M. F. Piehler. 2013a. Oyster-mediated benthic-pelagic coupling modifies nitrogen pools and processes. Marine Ecology Progress Series 493:23-30.

Smyth A.R., S.P. Thompson, K.N. Siporin, W.S. Gardner, and M.J. McCarthy. 2013b. Assessing nitrogen dynamics throughout the estuarine landscape. Estuaries and Coasts 36: 44-55.

Smyth, A. R., M. F. Piehler, and J. H. Grabowski. 2015. Habitat context influences nitrogen removal by restored oyster reefs. Journal of Applied Ecology 52:716-725.

Smyth, A.R., Geraldi, N.R., Thompson, S.P., Piehler, M.F. 2016. Biological activity exceeds biogenic structure in influencing sediment nitrogen cycling in experimental oyster reefs. Marine Ecology Progress Series 560: 173-183.

Testa, J. M., D. C. Brady, J. C. Cornwell, M. S. Owens, L. P. Sanford, C. R. Newell, S. E. Suttles, and R. I. E. Newell. 2015. Modeling the impact of floating oyster (Crassostrea virginica) aquaculture on sediment-water nutrient and oxygen fluxes. Aquaculture Environment Interactions 7:205-222.