



Land Use Methods and Metrics Outcome

Peter Claggett¹, Labeeb Ahmed¹, Jacob Czawlytko², Jarlath O'Neil-Dunne³, Sarah McDonald¹, Patrick McCabe², Sean MacFaden³, Rachel Soobitsky², and Renee Thompson¹

¹ Lower Mississippi-Gulf Water Science Center, U.S. Geological Survey, Annapolis, MD 21403

² Chesapeake Conservancy, Annapolis, MD 21403

³ Spatial Analysis Laboratory, University of Vermont, Burlington, VT 05405

Through the Chesapeake Bay Watershed Agreement, the Chesapeake Bay Program has committed to...




<https://blog.nature.org/science/2016/09/08/energy-sprawl-is-the-largest-driver-of-land-use-change-in-the-u-s/>

Goal: Conserve landscapes treasured by citizens in order to maintain water quality and habitat; sustain working forests, farms and maritime communities; and conserve lands of cultural, indigenous and community value.

Outcome: Assess and understand the impacts of land use change on watersheds, habitats, and communities at a scale relevant to county-level decision-makers.


Through the Chesapeake Bay Watershed Agreement, the Chesapeake Bay Program has committed to...



- 1. Measure rate of farmland, forest and wetland conversion, and the extent and rate of change in impervious surface coverage.*
- 2. Quantify the potential impacts of land conversion to water quality, healthy watersheds and communities.*
- 3. Launch a public awareness campaign to share this information with citizens, local governments, elected officials and stakeholders.*



What is our Expected and Actual Progress?





Learn

What have we learned in the last two years?



Successes and Challenges

Challenges:


- Automating the classification of land cover and land use
- Separating signal from noise when mapping change in land conditions
- Outdated inventory of non-tidal wetlands
- Communicating the complexities of high-resolution land change

Successes:

- CBP Partnership support for monitoring both land cover and land use change at high resolution
- Refinement of land cover mapping techniques (Univ. of Vermont)
- Development of high-res land use mapping techniques (USGS, Chesapeake Conservancy)
- Emerging understanding of land cover/use change and potential impacts to water quality and healthy watersheds

Status of the National Wetlands Inventory Inventory October 2020

Vintage of the NWI in the majority of
watershed is ~1980's



Chesapeake Bay Program Land Use Classification (55 classes)

1. Water (8)

1.1 Lentic

- 1.1.1 Estuary
- 1.1.2 Lakes & Ponds

1.2 Lotic

- 1.2.1 Streams
 - 1.2.1.1 Sunlit
 - 1.2.1.2 Shaded
 - 1.2.1.3 Culverted/ Buried
- 1.2.2 Ditches
 - 1.2.2.1 Sunlit
 - 1.2.2.2 Shaded
 - 1.2.2.3 Culverted/ Buried

2. Developed (12)

2.1 Impervious

- 2.1.1 Roads
- 2.1.2 Structures
- 2.1.3 Other Impervious (Parking lots, driveways)

2.2 Pervious

- 2.2.1 Turf Grass
- 2.2.2 Bare Developed
- 2.2.3 Suspended Succession (rights-of-way)
 - 2.2.3.1 Barren
 - 2.2.3.2 Herbaceous
 - 2.2.3.3 Scrub-shrub

2.3 Tree Canopy (TC)

- 2.3.1 TC over Roads
- 2.3.2 TC over Structures
- 2.3.3 TC over Other Impervious
- 2.3.4 TC over Turf Grass

3. Forest (6)

3.1 Forest (≥ 1 acre)

3.2 Harvested Forest

- 3.2.1 Barren
- 3.2.2 Herbaceous

3.3 Natural Succession (> 3 years)

- 3.3.1 Barren
- 3.3.2 Herbaceous
- 3.3.3 Scrub-shrub

4. Production (13)

4.1 Agriculture

- 4.1.1 Cropland
 - 4.1.1.1 Barren
 - 4.1.1.2 Herbaceous
- 4.1.2 Pasture
 - 4.1.2.1 Barren
 - 4.1.2.2 Herbaceous
- 4.1.3 Orchard/vineyard
 - 4.1.3.1 Barren
 - 4.1.3.2 Herbaceous
 - 4.1.3.3 Scrub-shrub

4.2 Solar fields

- 4.2.1 Barren
- 4.2.2 Herbaceous
- 4.2.3 Scrub-shrub
- 4.2.4 Impervious

4.3 Extractive

- 4.3.1 Barren
- 4.3.2 Impervious

5. Wetlands and Water Margins (16)

5.1 Tidal

- 5.1.1 Barren
- 5.1.2 Herbaceous
- 5.1.3 Scrub-shrub


5.2 Non-tidal

- 5.2.1 Riverine - Floodplain
 - 5.2.1.1 Barren
 - 5.2.1.2 Herbaceous
 - 5.2.1.3 Scrub-shrub
 - 5.2.1.4 Forest
- 5.2.2 Riverine - Headwater
 - 5.2.2.1 Barren
 - 5.2.2.2 Herbaceous
 - 5.2.2.3 Scrub-shrub
 - 5.2.2.4 Forest
- 5.2.3 Terrene
 - 5.2.3.1 Barren
 - 5.2.3.2 Herbaceous
 - 5.2.3.3 Scrub-shrub
 - 5.2.3.4 Forest


5.3 Bare shore

Enhanced Resolution Hydrography Lower Susquehanna Example

National Hydrography Dataset, 1:24,000
6,923.6 km



CBP Hyper-Resolution Flowpaths, 1:2000
16,784.6 km




Why the 2x difference in “stream” length?

Added
Tributaries

Increased
Complexity

Extended
headwaters



— NHD24K

— HyperRes



Tree Canopy Change in Two Suburban Counties

Prince George's County: 2014 - 2018

TC Loss (7,673 acres):

- 59% of loss change occurred within forest or wetlands
- 41% of loss occurred in developed areas

TC Gain (518 acres):

- 16% of gain occurred within forest or wetlands
 - shrub/scrub; edge of forest
- 54% of gain occurred in developed areas
- 29% of gain occurred on agricultural lands

Anne Arundel County: 2014 - 2018

TC Loss (2,544 acres):


- 57% of loss change occurred within forest or wetlands
- 42% of loss occurred in developed areas

TC Gain (188 acres):



- 9% of gain occurred within forest or wetlands
 - shrub/scrub; edge of forest
- 55% of gain occurred in developed areas
- 35% of gain occurred on agricultural lands

Parcel-Level Deconstruction of Urban Development (1985 – 2017)

Year-Built Attributes
from Tax Records



Year-Built Attributes
from USGS' LCMAP



Informing Land Use and Conservation Decisions with Data





On the Horizon

- Policies and regulations related to climate change with land use implications
- CBP Partner commitments to conserving 30% of land by 2030 and 50% by 2050
- Development of an effective CBP local engagement strategy (with science translators)
- Technological Innovations (e.g., LiDAR, satellite imagery, artificial intelligence)
- Increased understanding of the role of landscape context in estimating BMP efficiencies, pollutant loads, and impacts to stream flow and temperature
- CBP Partner uses and interest in the high-resolution land use/cover products



Adapt

How does all of this impact our work?



Based on what we learned, we plan to ...

- Monitor changes in land cover (12 classes), land use (55 classes), and watershed health metrics every four years: 2013-2017-2021.
- Deconstruct high-res land use from 2013 to mid 1980's
- Refine forecasts of land use change to include agriculture and timber harvests.
- Relate land use changes to effects on water quality, healthy watersheds, and communities.
- Formally publish the data
- Develop online viewers and other communication and interpretive products



Help

*How can the Management Board
lead the Program to adapt?*



Help Needed

- Support re-soliciting a Cooperative Agreement to monitor land use/cover change every 4-5 years through 2030 (e.g., adding 2025/26 and 2029/30 dates).
- Support integrating this outcome with the Land Use Options Evaluation and Local Leadership Outcomes, and with the Local Engagement Strategy to help ensure that data are used to inform land use planning and land conservation decisions.



Discussion